☆ THE ART OF COMPUTER PROGRAMMING ☆

EARLIEST ERRATA TO VOLUME 2 (3rd edition)

This document is a transcript of the notes that I had been making in my personal copy of *The Art of Computer Programming*, Volume 2 (third edition) since it was first printed in 1997.

Four levels of updates — "errors," "amendments," "plans," and "improvements" — appear, indicated by four different typographic conventions:

▸**Page 666** line 1 ——— 04 Jul 1776

Technical or typographical errors (aka bugs) are the most critical items, so they are flagged with a '▸' preceding the page number. The date on which I first was told about the bug is shown; this is the effective date on which I paid the finder's fee. The necessary corrections are indicated in a straightforward way. If, for example, the book says 'n' where it should have said '$n+1$', the change is shown thus:

$n \leadsto n+1$

Page 666 line 2 ——— 14 Jul 1789

Amendments to the text appear in the same format as bugs, but without the '▸'. These are things I wish I had known about or thought of when I wrote the original text, so I added them later. The date is the date I drafted the new text.

Page 666 line 3 . . . . . 20 Nov 1917

Plans for the future represent a third kind of item. In such notes I sketched my intentions about things that I wasn't ready to flesh out further when I wrote them down. You can identify these items because they're written in slanted type, and preceded by a bunch of dots '. . . . . ' leading to the date on which I recorded the plan in my files.

Page 666 line 4 ——— 10 Jan 1938

The fourth and final category — indicated by page and line number in smaller, slanted type — consists of minor corrections or improvements that most readers don't want to know about, because they are so trivial. You wouldn't even be seeing these items if you hadn't specifically chosen to print the complete errata list in all its gory details. Are you sure you wanted to do that?

My shelves at home are bursting with preprints and reprints of significant research results that I want to digest and summarize, where appropriate, in the ultimate edition of Volume 2. I didn't do that in the third edition because I would surely have to do it over again later: New results continue to pour forth at a great rate, and I will have time to rewrite that volume only once. Volumes 4 and 5 need to be finished first. So I've put most of my effort so far into writing up those parts of the total picture that seem to have converged to their near-final form. It follows, somewhat paradoxically, that the updates in this document are most current in the areas where there has been least activity.

On the other hand I do believe that the changes listed here bring Volume 2 completely up to date in two respects: (1) All of the research problems in the previous edition — i.e., all exercises that were rated 46 and above — have received new ratings of 45 or less whenever I learned of a solution; and in such cases, the answer now refers to that solution. (2) All of the historical information about pioneering developments has been amended whenever new details have come to my attention.

The ultimate, glorious, future editions of Volumes 1–3 are works in progress. Please let me know of any improvements that you think I ought to make. Send your comments either by snail mail to D. E. Knuth, Computer Science, Gates Building 1B, Stanford University, Stanford CA 94305-9015, or by email to `taocp@cs.stanford.edu`. *(Use email for book suggestions only, please — all other correspondence is returned unread to the sender, or discarded, because I have no time to read ordinary email.) Although I'm working full time on Volume 4 these days, I will try to reply to all such messages within a year of receipt. Current news about The Art of Computer Programming is posted on*

`http://www-cs-faculty.stanford.edu/~knuth/taocp.html`

and updated regularly.

— Don Knuth, September 1997

Oh! If only someone would give me time, time,
time to do everything properly,
to read everything at my own *tempo,*
to take it apart and put it together again.
— KARL BARTH (1922)

☆☆☆☆☆ SEMINUMERICAL ALGORITHMS ☆☆☆☆☆

Copyright © 1997, 1998, 1999, 2000, 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009, 2010, Addison–Wesley; all rights reserved
Last updated 16 August 2022

All of these corrections have been made in subsequent printings.

Page iv line 3 of the Library of Congress Data — 07 Jan 2011

`xiv,762p.` ↝ `xiv,764p.`

Page vii line 17 — 28 Oct 2008

cheerfully pay \$2.56 ↝ cheerfully award \$2.56

Page ix line 11 — 09 Oct 2010

This is sometimes unfortunate ↝ A motley mixture is, however, often unfortunate

Page x line −6 — 09 Oct 2010

This is not ↝ This distinction is not

Page xi line 9 — 09 Oct 2010

Later editions ↝ Later printings

▸**Page xii** line 11 ("General Test Procedures") — 22 Dec 1997

41 ↝ 42

▸**Page xii** line −10 — 19 Mar 2010

Multiple Precision Arithmetic ↝ Multiple-Precision Arithmetic

Page xiii replacement for the bottom line — 07 Jan 2011

Appendix C — Index to Algorithms and Theorems 735

Index and Glossary 737

Page 2 a new paragraph near the top — 27 Feb 2004

f) *Cryptography.* A source of unbiased bits is crucial for many types of secure communications, when data needs to be concealed.

[Now relabel the paragraphs about "aesthetics" and "recreation."]

Page 3 lines 11–13 — 05 Sep 2004

[See the articles by Kendall ... 61. See also S. H. Lavington's ↝ [F. N. David describes the early history in *Games, Gods, and Gambling* (1962). See also Kendall ... 61; S. H. Lavington's

Page 5 line 8 — 09 Oct 1998

$\lfloor X/10^9 \rfloor$, ↝ $\lfloor X/10^9 \rfloor$,

Page 9 exercise 19 — 08 May 2005

[*M48*] Solve the problems of exercises 11 through 15 ↝ [*HM47*] Solve the problems of exercises 11 through 15 asymptotically

Page 9 the rating of exercise 21 — 05 Apr 2007

[*42*] ↜ [*40*]

▸**Page 12** line −4 — 26 Oct 1998

(See exercise 3.) ↜

▸**Page 16** line 5 — 29 Dec 1997

do not to provide ↜ do not provide

▸**Page 17** line 21 — 03 Jan 2002

383–389 ↜ 163–167

Page 17 line −10 — 21 May 1998

then ↜ *then*

▸**Page 18** line 11 — 26 Jan 2001

length λ of ↜ length λ of the period of

Page 20 line 15 — 08 Nov 2005

called *the order of a modulo m* ↜ called the *order* of a modulo m

Page 24 between lines 1 and 2 — 13 Jan 1998

[there's a spurious dot near the left margin]

Page 26 new copy for exercise 8 — 27 Feb 2004

Let $Y_n = \lfloor 10X_n/2^{35} \rfloor$ ... much more often ↜
Let $Y_n = \lfloor 20X_n/2^{35} \rfloor$; then Y_n should be a random integer between 0 and 19, and the triples $(Y_{3n},\ Y_{3n+1},\ Y_{3n+2})$ should take on each of the 8000 possible values from $(0, 0, 0)$ to $(19, 19, 19)$ with nearly equal frequency. But with 1,000,000 values of n tested, many triples never occurred, and others occurred much more often

Page 28 top — 14 Nov 1997

[the page number is blurry]

Page 28 line 13 — 04 Oct 2007

$k \leftarrow 55$. ↜ $k \leftarrow 55$. (We cannot have both $j = 0$ and $k = 0$.)

▸**Page 29** line −15 — 04 Jan 1998

came as shock ↜ came as a shock

Page 31 line 8 — 18 Oct 2006

Section 7.1 ↜ Section 7.1.3

▸**Page 33** line −2 — 29 Dec 1997

from (12) ↜ from (13)

▸**Page 33** bottom line — 08 Dec 2009

222 ↜ 222–227

Page 39 lines 9 and 10 — 08 Jul 2001

Lemma 3.2.1.2Q, the trick of exercise 7 ↜ the results of exercises 7 and 13

Page 41 line 14 — 21 Jan 2009

Dr. I. J. Matrix ↝ Dr. I. J. Matrix and Donald C. Rehkopf

▸**Page 48** line 4 after Fig. 3 — 20 Nov 1997

random fracton ↝ random fraction

Page 49 in the caption to Fig. 4 — 11 Dec 2007

distributions. ↝ distributions. The x value marked "5%" is the percentage point where $F(x) = 0.05$.

Page 50 in (11) — 03 Feb 2005

max ↝ sup (twice)

Page 50 line 22 — 03 Feb 2005

maximum ↝ least upper bound

▸**Page 54** bottom line — 07 Feb 2000

every few microseconds ↝ every few milliseconds

▸**Page 69** line −2 — 30 Apr 2008

$-6C^{-1}$ ↝ $-6C_1^{-1}$

Page 69 bottom line — 11 Dec 2009

$V \approx Q^T C_1^{-1} Q/(n-6)$ ↝
by (10), $V \approx Q^T C_1^{-1} Q/(n-6)$, where $Q = (Q_1 \ldots Q_t)^T$.

▸**Page 72** line 5 — 22 Apr 2002

8 (1990) ↝ **9** (1990)

Page 74 line 19 — 22 Aug 2002

Paris **81** (1875) ↝ **81** (Paris, 1875)

▸**Page 74** line 21 — 03 Oct 2008

(*Proc. Royal* ↝ [*Proc. Royal*

▸**Page 78** exercise 24 — 03 Feb 2005

line 1: [*HM35*] ↝ [*HM37*]
line 7: $N(\alpha)$ ↝ $N(\alpha)^2$ (twice)

▸**Page 81** replacement for line −6 — 08 Nov 1997

$$\delta(x) = \lfloor x \rfloor + 1 - \lceil x \rceil = [x \text{ is an integer}]; \tag{6}$$

Page 82 the line following (10) — 13 Jun 2001

exercise 2 ↝ exercises 1.2.4–38 and 1.2.4–39(a,b,g)

▸**Page 89** line −13 — 25 Jul 2005

(1978) ↝ (1977)

▸**Page 90** line 4 — 25 Feb 1999

$\sum_{j=1}^{t} a_t$ ↝ $\sum_{j=1}^{t} a_j$

Page 90 new exercise — 13 Jun 2001

[Renumber exercise 3 as exercise 2, delete the former exercise 2, and insert the following new material:]

3. [*M23*] (N. J. Fine.) Prove that $|\sum_{k=0}^{n-1}((2^k x + \frac{1}{2}))| < 1$ for all real numbers x.

Page 92 exercise 22 — 03 Feb 2005

If x is a real number ↛ If x is a random real number, uniformly distributed

▸**Page 92** rating of exercise 23 — 28 Apr 1999

[*28*] ↛ [*M28*]

Page 103 lines 6–8 of step S7 — 25 Aug 1999

In hundreds ... dimensions. ↛ Usually $|z_j| \le 1$, but L. C. Killingbeck noticed in 1999 that larger values occur for about 0.00001 of all multipliers when $m = 2^{64}$.

Page 106 improvement to Table 1 — 02 May 2004

L. C. Killingbeck has carried out an exhaustive search of all multipliers $a \equiv 1 \bmod 4$ when $m = 2^{32}$. He found that the multiplier $a = 2650845021$ has $\nu_2^2 = 4938969760$, $\nu_3^2 = 2646962$, $\nu_4^2 = 68342$, $\nu_5^2 = 8778$, and $\nu_6^2 = 1506$, so it outperforms the other multipliers listed for this modulus. Indeed, its spectacular μ scores $(3.61, 4.20, 5.37, 8.85, 4.11)$ exceed all values of μ_2, μ_3, μ_4, and μ_5 for any modulus in the entire table.

▸**Page 107** line number 25 of the table — 20 Nov 1998

15.6 ↛ 14.6

Page 109 line −16 — 28 Sep 2002

p. 332 ↛ 332

Page 111 line 1 — 04 Nov 2001

(44) ↛ (44)

Page 112 in (49) — 06 Feb 2005

remove the factor $\dfrac{1}{N}$ (three times)

Page 113 near the top — 06 Feb 2005

remove the factors $1/N$ from (52), (53), and the line after (53); also change $0 < k < m$ ↛ $0 \le k < m$ on line 4

Page 113 line 2 of Theorem N — 06 Feb 2005

length m ↛ *length* $m > 1$

▸**Page 115** line 1 of exercise 8 — 16 Feb 1999

[*M16*] Line 18 ↛ [*M18*] Line 10

Page 117 exercise 26 — 06 Feb 2005

line 2: N^{-1} ↛
line 3: Where does ... $m = 1$? ↛

Page 117 last line of exercise 23 — 05 Jan 2001

Alekseev ↝ Alexeev

▸**Page 122** line −4 — 30 Sep 2003

0.587 ↝ 0.590

▸**Page 122** replacement for line −3 — 07 Feb 2000

P4. [Compute X_1, X_2.] If $S = 0$, set $X_1 \leftarrow X_2 \leftarrow 0$; otherwise set

Page 122 line −2 — 22 Dec 1998

$=$ ↝ $\leftarrow$ (twice)

▸**Page 124** replacement for Figure 9 — 07 Feb 2000



Page 125 lines 6 and 7 after the figure — 25 Feb 1998

concave downward, and when $x > 1$ it is concave upward, ↝ concave, and when $x > 1$ it is convex,

▸**Page 127** line −6 — 02 Dec 1999

$15 \le j \le 31$ ↝ $16 \le j \le 31$

▸**Page 128** in Table 1 — 28 Jul 2000

the value of Y_4: -33.13 ↝ -33.16
the value of Y_5: -39.55 ↝ -39.51
the value of Y_9: 0.00 ↝
the value of Z_4: 34.93 ↝ 34.96
the value of Z_5: 41.35 ↝ 41.31
the value of Z_9: 0.00 ↝

▸**Page 129** replacement for lines −3 and −2 — 30 Sep 2003

Otherwise set V to a new uniform deviate; and repeat step F4 if the new V is $\le U$. Otherwise (that is, if K is even, in the discussion above), replace U by a new uniform deviate $(.b_0 b_1 \ldots b_t)_2$ and go back to F3.

Page 132 new paragraph following line 15 — 21 Mar 2005

With an improved approximation to the acceptance region [see J. L. Leva, *ACM Trans. Math. Software* **18** (1992), 449–455] we can, in fact, reduce the expected number of logarithm computations to only 0.012.

▸**Page 135** line −16 — 23 Mar 1999

$(1 - 2\nu)$ ↯ $(1 - 2/\nu)$

Page 136 line 9 — 11 Dec 2009

consists only of integer values ↯ is nonzero only at integer values

Page 138 line 1 of exercise 3 — 07 Feb 2000

dividing by k ↯ computing its *remainder* mod k

Page 139 exercise 9 — 25 Feb 1998

concave downward for $x < 1$, concave upward for $x > 1$? ↯ concave for $x < 1$, convex for $x > 1$?

▸**Page 139** exercise 12 — 19 Oct 1999

$f(x) < f(y)$ ↯ $f(x) > f(y)$

Page 140 bottom line — 19 May 2005

$X_1 \vee (X_2 \wedge (X_3 \vee (X_4 \wedge X_5)))$ ↯ $X_1 \mid (X_2 \,\&\, (X_3 \mid (X_4 \,\&\, X_5)))$

Page 141 line 2 of exercise 27 — 18 Oct 2006

Section 7.1 ↯ Section 7.1.3

▸**Page 141** line 1 of exercise 29 — 10 Nov 1997

a simple ↯ Find a simple

Page 145 lines −13 and −12 — 20 Aug 2001

division by j should *not* be used to determine k ↯ k should *not* be computed by taking a remainder modulo j

▸**Page 146** line 14 — 17 Mar 2001

Algorithm S ↯ Algorithm P

▸**Page 147** line 1 of exercise 14 — 09 Mar 2001

to sequence ↯ to a sequence

Page 148 new exercise — 26 May 2010

▸**18.** [*M32*] People sometimes try to shuffle n items $(X_1, X_2, \ldots, X_n)$ by successively interchanging

$$X_1 \leftrightarrow X_{k_1},\ X_2 \leftrightarrow X_{k_2},\ \ldots,\ X_n \leftrightarrow X_{k_n},$$

where the indices k_j are independent and uniformly random between 1 and n.

Consider the directed graph with vertices $\{1, 2, \ldots, n\}$ and with arcs from j to k_j for $1 \le j \le n$. Describe the digraphs of this type for which, if we start with the elements $(X_1, X_2, \ldots, X_n) = (1, 2, \ldots, n)$, the stated interchanges produce the respective permutations (a) $(n, 1, 2, \ldots)$; (b) $(1, 2, \ldots, n)$; (c) $(2, \ldots, n, 1)$. Conclude that these three permutations are obtained with wildly different probabilities.

Page 148 another new exercise — 12 Jan 2009

▸**19.** [*M28*] (*Priority sampling.*) Consider a file of N items in which the kth item has a positive weight w_k. Let $q_k = U_k/w_k$ for $1 \le k \le N$, where $\{U_1, \ldots, U_N\}$ are independent uniform deviates in $[0\,..\,1)$. If r is any real number, define

$$\widehat{w}_k^{(r)} = \begin{cases} \max(w_k, 1/r), & \text{if } q_k < r; \\ 0, & \text{if } q_k \ge r; \end{cases} \qquad \widehat{w}_k^{(r+)} = \begin{cases} \max(w_k, 1/r), & \text{if } q_k \le r; \\ 0, & \text{if } q_k > r. \end{cases}$$

a) If r is the nth smallest element of $\{q_1, \ldots, q_N\}$, prove that the expected value $\mathrm{E}\,\widehat{w}_1^{(r)}\widehat{w}_2^{(r)}\ldots w_k^{(r)}$ is $w_1 w_2 \ldots w_k$, for $1 \le k < n \le N$. *Hint:* Show that, if s is the $(n-k)$th smallest element of $\{q_{k+1}, \ldots, q_N\}$, we have $\widehat{w}_1^{(r)}\ldots\widehat{w}_k^{(r)} = \widehat{w}_1^{(s+)}\ldots\widehat{w}_k^{(s+)}$. (Notice that the quantity s is independent of $\{U_1, \ldots, U_k\}$.)

b) Consequently $\mathrm{E}\,\widehat{w}_{j_1}^{(r)}\ldots\widehat{w}_{j_k}^{(r)} = w_{j_1}\ldots w_{j_k}$ when $j_1 < \cdots < j_k$.

c) Show that, if $n > 2$, the variance $\mathrm{Var}(\widehat{w}_{j_1}^{(r)} + \cdots + \widehat{w}_{j_k}^{(r)})$ is $\mathrm{Var}(\widehat{w}_{j_1}^{(r)}) + \cdots + \mathrm{Var}(\widehat{w}_{j_k}^{(r)})$.

d) Given n, explain how to modify the reservoir sampling method so that the value of r and the $n-1$ items with subscripts $\{j \mid q_j < r\}$ can be obtained with one pass through a file of unknown size N. *Hint:* Use a priority queue of size n.

Page 148 quotation to follow the exercises — 24 Mar 2006

By means of the thread one understands the ball of yarn,
so we'll be satisfied and assured by having this sample.

— MIGUEL DE CERVANTES, *El Ingenioso Hidalgo Don Quixote de la Mancha* (1605)

Page 150 and throughout this section — 25 Oct 2010

[Place '∎' after each **Definition**, to delimit definitions in the style used for theorems, algorithms, programs, etc. This change affects pages 150, 151, 155, 159, 161, 162, 163, 168, and 171.]

Page 155 line −2 — 22 Mar 2002

H. S. ↝ H. S.

Page 164 line 4 — 22 Aug 2002

Paris **A285** (1977) ↝ **A285** (Paris, 1977)

▸**Page 173** line 16 — 30 Nov 1997

0000, 0001, 1101, or 1110; ↝ 0000, 0011, 1101, or 1110;

Page 173 line −7 — 23 Jun 2001

Walsh transform ↝ Hadamard transform

Page 176 top — 14 Jan 2000

line 2: per year ↝ per Gregorian year
line 3: per second ↝ per Gregorian second

▸**Page 176** line −19 — 25 Jan 2003

$3R \approx 600000$ ↝ $\approx 3R = 600000$

▸**Page 176** line −16 — 20 Sep 2005

53.5 teraMIP-years ↝ 535 teraMIP-years

Page 177 lines 11–13 — 11 Jan 2000

normal ... 216. ↝ *entirely normal* to base b, and he stated Theorem C informally without apparently realizing that it required proof [*Rendiconti Circ. Mat. Palermo* **27** (1909), 247–271, §12.]

Page 179 lines 21–23 — 08 Aug 2005

Finally Levin . . . analyzing Algorithm L. ↝ Finally Charles Rackoff refined the methods of that paper by introducing and analyzing Algorithm L [see L. Levin, *J. Symbolic Logic* **58** (1993), 1102–1103].

Page 179 lines 25–26 — 20 Mar 2000

[*STOC* **21** (1989), 12–24; **22** (1990), 395–404] ↝ [*SICOMP* **28** (1999), 1364–1396]

Page 179 lines 27–28 — 27 Feb 2004

not surveyed here ... practical random number generation. ↝ beyond the scope of this book.

▸**Page 182** line 2 of exercise 31 — 26 Dec 2002

$U_n < \frac{1}{2}$ ↝ $U_j < \frac{1}{2}$

▸**Page 183** line 7 of exercise 42 — 26 Jan 2001

k-bit vectors, ↝ k-bit vectors, with $c \neq c'$,

Page 185 near the bottom — 18 Feb 2001

line −4: `(long)(MM/AA)` ↝ `MM / AA`
line −2: `(long)(X/QQ)` ↝ `(X/QQ)`

Page 186 near the top — 18 Feb 2001

line 8: `(long)(MMM/AAA)` ↝ `MMM / AAA`
line 10: `(long)(Y/QQQ)` ↝ `(Y/QQQ)`

Page 186 near the top — 05 Jul 2010

line 12: `Z+=MM;` ↝ `Z+=MM-1;`
line 14: is never zero; `Z` always ↝ always
line 15: and 2^{31} ↝ and $2^{31} - 1$, as recommended by Liviu Lalescu

Page 186 replacement for line −10 — 01 Jan 2002

```
void ran_array(long aa[],int n) {/* put n new values in aa */
```

Page 187 line 8 — 01 Jan 2002

[delete this line, since it is now redundant]

Page 187 replacement for lines 12–35 — 01 Jan 2002

```
register long ss=(seed+2)&(MM-2);
for (j=0;j<KK;j++) {
  x[j]=ss;                          /* bootstrap the buffer */
  ss<<=1; if (ss>=MM) ss-=MM-2;     /* cyclic shift 29 bits */
}
```

```
  x[1]++;                  /* make x[1] (and only x[1]) odd */
  for (ss=seed&(MM-1),t=TT-1; t; ) {
    for (j=KK-1;j>0;j--)
      x[j+j]=x[j], x[j+j-1]=0;                  /* "square" */
    for (j=KK+KK-2;j>=KK;j--)
      x[j-(KK-LL)]=mod_diff(x[j-(KK-LL)],x[j]),
      x[j-KK]=mod_diff(x[j-KK],x[j]);
    if (is_odd(ss)) {                    /* "multiply by z" */
      for (j=KK;j>0;j--)  x[j]=x[j-1];
      x[0]=x[KK];            /* shift the buffer cyclically */
      x[LL]=mod_diff(x[LL],x[KK]);
    }
    if (ss) ss>>=1; else t--;
  }
  for (j=0;j<LL;j++) ran_x[j+KK-LL]=x[j];
  for (;j<KK;j++) ran_x[j-LL]=x[j];
  for (j=0;j<10;j++) ran_array(x,KK+KK-1);    /* warm it up */
}
```

(This program incorporates improvements to the author's original *ran_start* routine, recommended by Richard Brent and Pedro Gimeno in November 2001.)

Page 188 line 1 — 26 Jan 2001

different sets ↝ disjoint sets

▸**Page 188** line 17 — 21 Feb 2001

`void main()` ↝ `int main()`

Page 188 line 23 — 17 Sep 2000

`ran_x[0]);` ↝ `ran_x[0]); return 0;`

Page 188 line 25 — 01 Jan 2002

461390032 ↝ 995235265

▸**Page 189** line −15 — 12 Nov 2007

125–136 ↝ 125–137

Page 191 line −3 — 31 May 1999

in a shot ↝ by a shot

Page 193 line 4 of exercise 14 — 04 Dec 2000

(a,200) ↝ (a, 200)

Page 195 replacement for line −13 — 19 Nov 2008

either `0`, `1`, `2`, `3`, `4`, `5`, `6`, `7`, `8`, `9`, `a`, `b`, `c`, `d`, `e`, `f`
or `0`, `1`, `2`, `3`, `4`, `5`, `6`, `7`, `8`, `9`, `A`, `B`, `C`, `D`, `E`, `F`.

Page 196 line 2 (not an error) — 05 Jun 2008

[The spelling 'archæologists' here is intentionally archaic.]

▸**Page 196** line −11 — 02 Feb 1998

1942), 37–62; ↜ 1941), 37–67;

Page 196 lines −11 and −10 — 20 Feb 2003

"Ancient Mesoamerican computing practices," ... to appear. ↜ "Pratiche del calcolo nell'antica mesoamerica," *Storia della Scienza* **2** (Rome: Istituto della Enciclopedia Italiana, 2001), 976–990.

Page 198 top line — 17 Oct 2007

A.D. 501 ↜ A.D. 500 or 501

Page 198 line 22 — 06 Apr 2006

1596–1610 ↜ 1586–1610

Page 198 replacement for lines 23–33 — 01 Nov 2001

Decimal fractions began to appear sporadically in Europe; for example, a so-called "Turkish method" was used to compute $153.5 \times 16.25 = 2494.375$. Giovanni Bianchini developed them further, with applications to surveying, prior to 1450; but like al-Uqlīdisī, his work seems to have had little influence. Christof Rudolff and François Viète suggested the idea again in 1525 and 1579. Finally, an arithmetic text by Simon Stevin, who independently hit on the idea of decimal fractions in 1585, became popular. Stevin's work, and the discovery of logarithms soon afterwards, made decimal fractions commonplace in Europe during the 17th century. [For further remarks and references, see D. E. Smith, *History of Mathematics* **2** (1925), 228–247; V. J. Katz, *A History of Mathematics* (1993), 225–228, 345–348; and G. Rosińska, *Quart. J. Hist. Sci. Tech.* **40** (1995), 17–32.]

▸**Page 199** line 6 — 03 Dec 1998

de Seuil ↜ du Seuil

▸**Page 199** line 9 — 02 Nov 2002

people would ↜ people

Page 199 line 15 — 10 Nov 2002

(London, 1687) ↜ (London: 1687)

Page 199 bottom line — 27 Aug 2004

de Caramuel Lobkowitz ↜ Caramuel de Lobkowitz

Page 200 line 5 — 22 Aug 2002

(Paris: 1703) ↜ (Paris, 1703)

▸**Page 201** line 16 — 25 Oct 1999

Colenne ↜ Collenne

Page 202 line −16 — 22 Mar 2002

CACM **2** ↜ *CACM* **2**, 12

Page 202 line −3 — 10 Aug 1998

signed-magnitude ↝ *signed magnitude*

Page 203 lines 21 and 22 — 14 Oct 1999

the largest negative number representable in p digits is $500\ldots0$, and it is not ↝ the p-digit negative number $500\ldots0$ is not

Page 203 line −5 — 10 Aug 1998

signed- ↝ signed

▸**Page 205** line 8 — 06 Nov 1997

Marcziński ↝ Marczyński

▸**Page 205** line 10 — 26 Sep 2005

274–276 ↝ 274–277

Page 205 line 18 — 26 Sep 2005

245–247. ↝ 245–247; **4** (1961), 355.

Page 208 line 5 — 08 Apr 2000

$\bar{1}\ 1\ 0\ 1\ 0$ ↝ $\bar{1}\ 1\ 0\ 1$

Page 208 line 17 and lines 23–24 — 22 Aug 2002

Paris **11** (1840) ↝ **11** (Paris, 1840)

Page 208 line 24 — 07 Dec 2004

mechanical devices for arithmetic. ↝
mechanical devices for arithmetic. Thomas Fowler independently invented and constructed a balanced ternary calculator at about the same time [see *Report British Assoc. Adv. Sci.* **10** (1840), 55; **11** (1841), 39–40].

Page 209 line 5 — 23 Mar 2006

this system, ↝ this system, which was known in 13th-century India,

Page 209 line 8 — 07 Feb 2006

Algorithm 3.3.2P ↝ Algorithms 3.3.2P and 3.4.2P

▸**Page 210** line 1 — 26 Jan 2001

number 1). ↝ number 1.)

Page 210 line 2 of exercise 6 — 10 Aug 1998

signed-magnitude ↝ signed magnitude

▸**Page 210** line 2 of exercise 12 — 02 Dec 1999

$(b_{n+1}\ldots b_0)_{-2}$ ↝ $(b_{n+2}\ldots b_0)_{-2}$

Page 214 lines 19, 20, −8, −7 — 28 Jan 1999

2252 ↝ 2214 (twice)
6256 ↝ 6261 (twice)

Page 214 and following . . . . . 16 Apr 1999

In the fourth edition I will change the definitions so that we have $|f| < b$; equation (2) will become "$-b^p < b^{p-1}f < b^p$" and (5) will become "$1 \le |f| < b$"; Avogadro's number with an excess-50 decimal representation will be $(73, +6.0221400)$, etc. The net effect will be to change the notion of excess, so that what was excess q in the third edition will be called excess $q+1$ in the fourth edition. This change in terminology will work much better with respect to IEEE standard floating point conventions (and it will properly match the term "ulp," defined at the bottom of page 232), although equations (21)–(24) will become slightly more messy.

Page 216 in Fig. 2 — 11 Feb 2007

[On the branch from A4 to A7, change the label '$e_u \gg e_v$' to '$e_u \ge e_v + p + 2$']

▸**Page 219** replacement for program lines 26–28 — 10 Nov 2002

```
26      LD1   FV(0:1)   Check for opposite signs.
27      JAP   1F
28      J1N   N2        If not, normalize.
29      JMP   2F
30  1H  J1P   N2
31  2H  SRC   5         |rX| ↔ |rA|.
```

[Now lines 29–67 must be renumbered 32–70. Also change '4' to '5' on (new) line 38, 'N3A' to 'N2' on (new) line 39, and delete 'N3A' from (new) line 47.]

▸**Page 219** near the bottom — 10 Nov 2002

line −6: 25 to 37 ↛ 26 to 40
line −1: 55 uses ↛ 58 uses

▸**Page 221** line −8 — 10 Nov 2002

30–34 ↛ 33–37

Page 228 line 3 of exercise 16 — 04 Oct 2010

and d ↛ and d, with $c + di \neq 0$

Page 232 lines 7 and 8 — 30 Nov 2000

(15) ↛ (15) and (16) ↛ (16)

Page 232 line 15 — 01 Mar 2007

$b^{e-1} \le x < b^e$ ↛ $b^{e-1} \le |x| < b^e$

Page 238 near the bottom — 28 Jan 1999

lines −9 and −1: 60225 ↛ 60221
lines −9 and −8: 66256 ↛ 66261
line −8: 2252 ↛ 2214

▸**Page 239** bottom line — 20 Mar 1999

$\delta_{(u\oplus v)\oplus w)}$ ↛ $\delta_{(u\oplus v)\oplus w}$

▸**Page 240** line −4 — 25 Mar 2001

$[u_0 \mathbin{▽} v_0, u_1 \mathbin{△} v_1] \leadsto [u_0 \mathbin{▽} v_0 \,..\, u_1 \mathbin{△} v_1]$

Page 240 replacement for bottom two lines — 28 Jan 1999

$$N = \big[(24, +.60221331) \,..\, (24, +.60221403)\big],$$
$$h = \big[(-26, +.66260715) \,..\, (-26, +.66260795)\big];$$

Page 241 replacement for lines 2 and 3 — 28 Jan 1999

$$N \oplus h = \big[(24, +.60221331) \,..\, (24, +.60221404)\big],$$
$$N \otimes h = \big[(-2, +.39903084) \,..\, (-2, +.39903181)\big].$$

Page 243 replacement for lines 15–17 — 28 Jan 1999

c) $u = (24, +.60221400)$, $v = (27, +.00060221)$;
d) $u = (24, +.60221400)$, $v = (31, +.00000006)$;
e) $u = (24, +.60221400)$, $v = (28, +.00000000)$.

▸**Page 244** line 8 of exercise 19 — 10 Mar 2007

$s_n = \leadsto s_n - c_n =$

Page 245 line 3 — 21 Jul 2007

$-0 < +0 \leadsto -0 < 0 < +0$

Page 246 line 21 — 16 Jul 2006

denormal numbers $\leadsto$ *subnormal numbers*

▸**Page 246** lines 21 and 22 — 02 Nov 2004

and the so-called NaNs that represent infinite, undefined, or otherwise unusual quantities $\leadsto$ *infinities, and the so-called NaNs that represent undefined or otherwise unusual quantities*

Page 247 line 8 — 24 Sep 1999

as much more time $\leadsto$ as much time

Page 247 line −11 — 10 Aug 1998

signed-magnitude $\leadsto$ signed magnitude

Page 255 line 19 — 19 Sep 2005

from different sources $\leadsto$ from many different sources

Page 262 line −7 — 18 Sep 2005

347. $\leadsto$ 347. See also A. Berger, L. A. Bunimovich, and T. P. Hill, *Trans. Amer. Math. Soc.* **357** (2004), 197–219.

Page 266 line 21 — 10 Aug 1998

signed-magnitude $\leadsto$ signed magnitude

▸**Page 267** line 4 — 26 Dec 2000

To A3 if $j = n$. $\leadsto$ Exit the loop if $j = n$.

▸**Page 268** line 6 — 10 Aug 2000

`STA  W,1` ↜↝ `STA  W+N,1`

▸**Page 268** line 24 — 08 Oct 2006

$(u_{m-1}\ldots u_1u_0)\times v_j$ ↜↝ $(u_{m-1}\ldots u_1u_0)_b\times v_j$

Page 268 line 26 — 22 Oct 2006

it is best ↜↝ it is simpler

Page 268 line −16 — 08 Apr 2002

$w_{m-2}\ \ldots,\ w_0$ ↜↝ $w_{m-2},\ \ldots,\ w_0$

Page 269 line 2 — 22 Oct 2002

84. ↜↝ 84

Page 269 line −18 — 08 Apr 2002

Program M. ↜↝ **Program M**

▸**Page 272** lines 10 and 23 — 23 Jan 1998

$\lfloor(b-1)/v_{n-1}\rfloor$ ↜↝ $\lfloor b/(v_{n-1}+1)\rfloor$

▸**Page 272** line 2 of step D3 — 03 Feb 2005

test if $\hat{q}=b$ ↜↝ test if $\hat{q}\geq b$

Page 272 line −3 — 07 Feb 2000

$\hat{q}\,(v_{n-1}\ldots v_1v_0)_b$ ↜↝ $\hat{q}\,(0\,v_{n-1}\ldots v_1v_0)_b$

Page 273 in steps D4 and D6 — 29 Dec 1997

$u_{n+j}u_{j+n-1}$ ↜↝ $u_{j+n}u_{j+n-1}$

Page 273 line 3 of step D6 — 01 Apr 2000

debugging. ↜↝ debugging. See exercise 22.

▸**Page 273** program line *041* — 20 Oct 1998

q ↜↝ $\hat{q}$

▸**Page 273** program line *044* — 03 Feb 2005

quotient $=b$ ↜↝ quotient $\geq b$

▸**Page 273** program line *046* — 22 Jan 1998

u_{j+n-11} ↜↝ u_{j+n-1}

▸**Page 274** line 2 — 19 Mar 1998

`U+B-1,2` ↜↝ `U+N-1,2`

▸**Page 275** line 19 — 23 Jan 1998

106 cycles ↜↝ 111 cycles

Page 275 line −4 — 10 Aug 1998

signed-magnitude ↜↝ signed magnitude

Page 279 line −14 — 28 Oct 2000

273–283. ↜ 273–283, **24** (1998), 359–367.

Page 279 line −10 — 22 Aug 2002

Paris (1719) ↜ (Paris, 1719)

Page 279 line −3 (*not* an error!) — 06 Jan 2010

["J. C. V. Hoffmann" is printed for Immanuel C. V. Hoffmann in the source.]

▸**Page 280** line 26 — 23 Apr 1998

millenium ↜ millennium

▸**Page 282** line 1 of exercise 16 — 17 Dec 1998

a algorithm ↜ an algorithm

▸**Page 282** line 2 of exercise 20 — 03 Feb 2005

u_{n-2}, then ↜ u_{n-2} and $\hat{q} < b$, then

▸**Page 282** replacement for line 2 of exercise 23 — 23 Jan 1998

have $\lfloor b/2 \rfloor \le v\lfloor b/(v+1) \rfloor < (v+1)\lfloor b/(v+1) \rfloor \le b$.

▸**Page 285** line −7 — 15 Nov 2001

of the Section 4.3.1 ↜ of Section 4.3.1

▸**Page 287** line 8 — 27 Oct 2004

Civilization ↜ *Civilisation*

Page 292 line 17 — 12 Jul 2003

its Applications ↜ *Its Applications*

Page 293 new sentence at end of exercise 5 — 05 Dec 2000

What is the largest possible $m_1 m_2 \ldots m_r$ when each residue u_j must fit in eight bits of memory?

▸**Page 301** line 23 — 30 Apr 2008

multiplications of p-bit numbers ↜ multiplications of q-bit numbers

▸**Page 303** in (25) — 07 Feb 1999

$O(n^{1.63})$ ↜ $O(n^{1.631})$

▸**Page 304** line 20 — 07 Feb 2000

positive integers e and f ↜ relatively prime positive integers e and f

Page 308 lines −17 *and* −16 — 07 Sep 2001

headed $\hat{u}_s$ in Table 1. The $\hat{v}_s$ column ↜ headed $2^7\,\hat{u}_s$ in Table 1. The column for $\hat{v}_s$

▸**Page 309** line 1 — 14 Aug 2005

$\omega_r = x_r + iy_r$ ↜ $\omega_r = x_r + iy_r$ and $r \ge 2$

▸**Page 310** replacement for line 15 — 08 Nov 1997

$$\left|(\hat{\hat{w}}_r)' - \hat{\hat{w}}_r\right| < 2^k(4k2^k - 2k)\delta + (2^k - 1)2k\delta; \quad |w'_r - w_r| < 4k2^k\delta. \qquad (50)$$

▸**Page 310** renumbering the equations — 08 Nov 1997

line 18: (50) ↛ (51)
lines 20 and 21: (51) ↛ (52)

▸**Page 310** line −12 — 27 Apr 1999

$m = 44$ ↛ $m = 144$

▸**Page 311** line 8 — 08 Nov 1997

(52) ↛ (53)

▸**Page 311** line 9 — 29 Jan 1998

$\lg \ldots \lg n \le 1$ ↛ $\lg \ldots \lg n \le 2$

▸**Page 311** line −9 — 29 Dec 1997

could than be done ↛ could then be done

▸**Page 313** line 13 — 15 May 1998

constant factor ↛ constant factor.

▸**Page 316** line 2 of exercise 5 — 14 Sep 2000

(21) can be ↛ (20) can be

Page 322 line −14 — 18 Oct 2006

Section 7.1 ↛ Section 7.1.3

▸**Page 324** line −7 — 10 Nov 2002

CE-11 ↛ EC-11

▸**Page 325** line 23 — 25 Oct 1999

.9 5 2 4 6 ↛ .9 5 2 6 4

▸**Page 328** replacement for exercise 9 — 14 May 2006

▸**9.** [*M29*] The purpose of this exercise is to compute $\lfloor u/10 \rfloor$ with binary shifting and addition operations only, when u is a nonnegative integer. Let $v_0(u) = 3\lfloor u/2 \rfloor + 3$ and

$$v_{k+1}(u) \;=\; v_k(u) + \lfloor v_k(u)/2^{2^{k+2}} \rfloor \qquad \text{for } k \ge 0.$$

Given k, what is the smallest nonnegative integer u such that $\lfloor v_k[u]/16 \rfloor \ne \lfloor u/10 \rfloor$?

Page 329 in exercise 19 — 19 May 2005

line 3: $U \leftarrow U - c_i(U \wedge m_i)$ ↛ $U \leftarrow U - c_i(U \,\&\, m_i)$
line 4: where "∧" denotes ↛ where "&" denotes

Page 330 quotation to follow line 8 — 11 Jun 2005

Irrationality is the square root of all evil.
— DOUGLAS HOFSTADTER, *Metamagical Themas* (1983)

Page 332 line 22 — 06 Jul 2006

29–47 ↝ 29–38, 39–47

Page 335 line -6 — 02 Dec 1999

and let F be a divisor ↝ and suppose that F is a divisor

Page 343 line 13 — 18 May 2002

Aryabhatiya (A.D. 499) by Aryabhata ↝ *Āryabhaṭīya* (A.D. 499) by Āryabhaṭa

Page 343 line 15 — 18 May 2002

Bháscara I in the sixth century ↝ Bhāskara I in the seventh century

▸**Page 345** line 11 — 14 Sep 1999

integer variables only ↝ integer values only

Page 348 line 12 — 14 Feb 2009

of the Algorithm B's behavior ↝ of Algorithm B's behavior

Page 348 line −15 — 23 Nov 1998

has discovered ↝ discovered

Page 349 in (36) — 11 Apr 1999

$0 \le x \le 1$ ↝ $0 < x \le 1$

Page 352 line 2 after (60) — 23 Nov 1998

[work in progress] has ↝

Page 352 line 2 after (61) — 23 Nov 1998

conjecture. ↝ conjecture. Vallée has successfully analyzed Algorithm B using rigorous "dynamical" methods of great interest [see *Algorithmica* **22** (1998), 660–685].

Page 355 the rating of exercise 32 — 23 May 2008

HM47 ↝ *HM42*

Page 357 line 8 — 18 May 2002

$(x_{n-1}$ ↝ $/(x_{n-1}$

Page 360 middle — 22 Aug 2002

line 17: *Paris* **11** (1733) ↝ **11** (Paris, 1733)
lines 25–26: *Paris* **19** (1844) ↝ **19** (Paris, 1844)

▸**Page 360** line −16 — 17 Jun 1998

a interesting ↝ an interesting

Page 360 lines −12 and −13 — 18 May 2002

[The statement of Corollary L should be in slanted type.]

▸**Page 362** in equation (25) — 30 Mar 2000

$\zeta(5)x^5$ ↝ $\zeta(5)x^4$

Page 366 lines −3 and −2 — 23 Nov 1998

Lecture Notes ... to appear. ↝ *Combinatorics, Probability and Computing* **6** (1997), 397–433; Flajolet and Vallée, *Theoretical Comp. Sci.* **194** (1998), 1–34.

▸**Page 368** replacement for lines 1–3 — 18 Feb 2004

Theorem E. *Let $p_k(a,n)$ be the probability that the $(k+1)$st quotient A_{k+1} in Euclid's algorithm is equal to a, when $u=n$ and when v is equally likely to be any of the numbers $\{0,1,\ldots,n-1\}$. Then*

Page 368 lines 15–18 — 07 Feb 2000

$= \leftsquigarrow \approx$ (four times)

Page 369 lines 12–13 — 07 Feb 2000

$= \leftsquigarrow \approx$ (three times)

▸**Page 370** line 3 — 05 Feb 1998

55.6 ↝ 5.6

Page 372 replacement for line 3 — 13 Jan 2000

$$\frac{1}{N^2}\sum_{m=1}^{N}\sum_{n=1}^{N} T(m,n) \;=\; \frac{2}{N^2}\sum_{n=1}^{N} nT_n - \frac{1}{2} - \frac{1}{2N}. \tag{57}$$

Page 372 line 14 — 13 Jan 2000

John D. Dixon and Hans A. Heilbronn. Dixon ↝ Gustav Lochs, John D. Dixon, and Hans A. Heilbronn. Lochs [*Monatshefte für Math.* **65** (1961), 27–52] derived a formula equivalent to the fact that (57) equals $(12\pi^{-2}\ln 2)\ln N + a + O(N^{-1/2})$, where $a \approx 0.065$. Unfortunately his paper remained essentially unknown for many years, perhaps because it computed only an average value from which we cannot derive definite information about T_n for any particular n. Dixon

Page 373 line 1 — 13 Jan 2000

to prove ↝ to confirm Lochs's work, proving

Page 373 line 3 — 07 Feb 2000

$\approx \leftsquigarrow =$

Page 373 new copy to follow (62) — 21 May 2005

D. Hensley proved in *J. Number Theory* **49** (1994), 142–182, that the variance of τ_n is proportional to $\log n$.

Page 375 lines 2 and 3 of exercise 13 — 30 Sep 2009

a polynomial with integer coefficients, having no rational roots, and having exactly one real root $\xi > 1$. ↝ a polynomial having exactly one real root $\xi > 1$, where ξ is irrational and $f'(\xi) \neq 0$.

▸**Page 377** line 8 of exercise 34 — 27 Feb 2004

$\sum_{y=1}^{n}$ ↝ $\sum$

Page 383 line 3 — 28 Sep 2002

$F(\alpha) = 1 \leadsto F(\alpha) = 1,$

Page 383 lines −3 and −2 — 07 Feb 2000

$= \leadsto \approx$ (two times)

▸**Page 384** lines 2 and 3 after (9) — 09 Jan 2009

P. Erdős and M. Kac, *Amer. J. Math.* **26** ↝ P. Erdös and M. Kac, *Amer. J. Math.* **62**

Page 384 lines −10 through −8 — 28 Dec 2000

[See D. E. Knuth ... 57–61.] ↝ [See D. E. Knuth, *Selected Papers on Analysis of Algorithms* (2000), 329–330, 336–337, for references to the relevant literature.]

Page 385 line −13 — 14 May 1998

mod p. ↝ mod p; see Arney and Bender, *Pacific J. Math.* **103** (1982), 269–294.

Page 386 replacement for lines 19–24 — 28 Dec 2000

Experiments by Tomás Oliveira e Silva indicate that $m(p)$ has an average value of about $2\sqrt{p}$, and it never exceeds $16\sqrt{p}$ when $p < 1000000000$. The maximum $m(p)$ for $p < 10^9$ is $m(850112303) = 416784$; and the maximum of $m(p)/\sqrt{p}$ occurs when $p = 695361131$, $m(p) = 406244$. According to these experimental results, almost all 18-digit numbers can be factored in fewer than 64,000 iterations of Algorithm B (compared to roughly 50,000,000 divisions in Algorithm A).

Page 387 line 3 — 31 Dec 2005

357.] ↝ 357. An equivalent idea had in fact been used already by Nārāyaṇa Paṇḍita in his remarkable book *Gaṇita Kaumudī* (1356); see Parmanand Singh, *Gaṇita Bhāratī* **22** (2000), 72–74.]

Page 387 line 10 — 07 Feb 2000

division: ↝ *multiplication or division:*

Page 387 Algorithm C — 12 Jun 2004

[I've changed the names of variables x and y in this algorithm to a and b, in order to avoid confusion with the x and y of (12) and (13).]

▸**Page 387** step C5 — 29 Dec 1997

Return to step C3 ↝ Return to step C4

Page 388 bottom line — 19 Sep 2005

solution x. ↝ solution x. Individual primes p and q are, however, preferable to moduli like p^2 unless p is quite small, because we tend to get even more information mod pq.

Page 390 line 12 — 19 May 2005

$\mathrm{rA} \leftarrow \mathrm{rA} \wedge S'[2, \mathrm{rI1}]$ ↝ $\mathrm{rA} \leftarrow \mathrm{rA}\ \&\ S'[2, \mathrm{rI1}]$

Page 392 line −4 — 20 Jul 2000

is composite ↜ cannot be prime

Page 394 line −3 — 20 Jun 2000

is prime ↜ is nonprime

Page 395 better wording of step P3 — 24 Jan 2003

If $j = 0$ ... step P5. ↜ If $y = n - 1$, or if $y = 1$ and $j = 0$, terminate the algorithm and say "n is probably prime." If $y = 1$ and $j > 0$, go to step P5.

▸**Page 395** near the bottom — 18 Dec 1998

line −10: about n ↜ about its input
line −9: certified a billion different primes ↜ tested a billion different numbers

Page 395 line −6 — 07 Feb 2000

cosmic radiations ↜ cosmic radiation

Page 396 upper middle — 19 Mar 2005

line 11: first proposed ↜ proposed
line 16: 35–36]. ↜ 35–36], and independently discovered by J. L. Selfridge.

Page 396 middle — 28 Apr 2005

A completely different ... purely of theoretical interest. ↜

A completely rigorous and deterministic way to test for primality in polynomial time was finally discovered in 2002 by Manindra Agrawal, Neeraj Kayal, and Nitin Saxena, who proved the following result:

Theorem A. *Let r be an integer such that $n \perp r$ and the order of n modulo r exceeds $(\lg n)^2$. Then n is prime if and only if the polynomial congruence*

$$(z + a)^n \equiv z^n + a \qquad (\text{modulo } z^r - 1 \text{ and } n)$$

holds for $0 \le z \le \sqrt{r}\,\lg n$. (See exercise 3.2.2–11(a).) ∎

An excellent exposition of this theorem has been prepared by Andrew Granville [*Bull. Amer. Math. Soc.* **42** (2005), 3–38], who presents an elementary proof that it yields a primality test with running time $\Omega(\log n)^6$ and $O(\log n)^{11}$. He also explains a subsequent improvement due to H. Lenstra and C. Pomerance, who showed that the running time can be reduced to $O(\log n)^{6+\epsilon}$ if the polynomial $z^r - 1$ is replaced by a more general family of polynomials. And he discusses refinements by P. Berrizbeitia, Q. Cheng, P. Mihăilescu, R. Avanzi, and D. Bernstein, leading to a probabilistic algorithm by which a proof of primality can almost surely be found in $O(\log n)^{4+\epsilon}$ steps whenever n is prime.

Page 398 replacement for lines 5 and 6 (see the following change) — 02 Jul 2001

After E1: 876 73 12 5329 1 —

Page 398 lines 1–5 of step E1 — 02 Jul 2001

$R' \leftarrow 2R$, ..., $n \bmod 2$. ↜ $R' \leftarrow 2R$, $U' \leftarrow R'$, $V \leftarrow D - R^2$, $V' \leftarrow 1$, $A \leftarrow \lfloor R'/V \rfloor$, $U \leftarrow R' - (R' \bmod V)$, $P' \leftarrow R$, $P \leftarrow (AR + 1) \bmod N$, $S \leftarrow 1$. (This algorithm follows

the general procedure of exercise 4.5.3–12, finding the continued fraction expansion of $\sqrt{kN}$. The variables U, U', V, V', P, P', A, and S represent, respectively, what that exercise calls $\lfloor\sqrt{D}\rfloor + U_n$, $\lfloor\sqrt{D}\rfloor + U_{n-1}$, V_n, V_{n-1}, $p_n \bmod N$, $p_{n-1} \bmod N$, A_n, and $n \bmod 2$, where n is initially 1.

Page 402 line 4 of the final paragraph — 08 Oct 2010

Annals of Math. ↝ *Annals of Math.* (2)

Page 402 bottom line — 14 May 1998

factors. ↝ factors. An excellent exposition of this method has been given by Joseph H. Silverman and John Tate in *Rational Points on Elliptic Curves* (New York: Springer, 1992), Chapter 4.

Page 403 line −19 — 07 Jul 1999

67 (1998), to appear ↝ **68** (1999), 429–451

Page 403 line −9 — 20 Jun 2000

to appear ↝ *Math. Comp.* **69** (2000), 1297–1304

Page 404 line −12 — 07 Feb 2000

take cube roots ↝ take cube roots quickly

Page 406 line 5 — 07 Jul 1999

considered ↝ published

Page 407 after line 10 — 16 Nov 2010

Historical note: It was revealed in 1997 that Clifford Cocks had considered the encoding of messages by the transformation $x^{pq} \bmod pq$ already in 1973, but his work was kept secret.

Page 409 line 10 — 26 Aug 1999

Joel Armengaud ↝ Joël Armengaud

Page 409 update on Mersenne primes . . . . . 02 May 2004

Tens of thousands of people have joined the GIMPS project, and Scott Kurowski has provided Internet software called PrimeNet to make the exponent assignments more systematic. Already in February 1998, more than one year's worth of computing was being done each day looking for Mersenne primes, and the amount of activity was growing steadily. Thus we can expect gradual progress on the hunt for these historic numbers.

I will list all Mersenne primes found after 1997 here, but on page 409 of the printed books I'll abbreviate the following information because space is tight. I won't update page 409 with every new printing, but I did make major amendments when preparing the 15th printing, and a few later ones (because I keep thinking mistakenly that the rate of discovery is about to cool down!). So far the new primes $2^p - 1$ found are:

p	*Discoverer*	*Date found*

3021377	*Roland Clarkson*	*27 January 1998*
6972593	*Nayan Hajratwala*	*1 June 1999*
13466917	*Michael Cameron*	*14 November 2001*
20996011	*Michael Shafer*	*17 November 2003*
24036583	*Josh Findley*	*May 2004*
25964951	*Martin Nowak*	*Feb 2005*
30402457	*Curtis Cooper, Steven Boone*	*Dec 2005*
32582657	*Curtis Cooper, Steven Boone*	*4 Sep 2006*
37156667	*Hans-Michael Elvenich*	*6 September 2008*
42643801	*Odd Magnar Strindmo*	*12 April 2009*
43112609	*Edson Smith*	*23 August 2008*

Page 409 line 4 before Theorem L — 08 Oct 2010

Annals of Math. ↛ *Annals of Math.* (2)

Page 412 near the top — 08 Jan 2005

lines 2 and 3: are prime [see ... 136]. ↛ are prime.
line 6: other forms. ↛ other forms. [See E. Picutti, *Historia Math.* **16** (1989), 123–136; Hugh C. Williams, *Édouard Lucas and Primality Testing* (1998), Chapter 2.]

Page 412 line 1 of exercise 11 — 20 Apr 1998

197209, $k = 5$, $m = 1$ ↛ 197209, $k = 5$, $m = 1$

Page 413 line 2 of exercise 22 — 21 Apr 2002

given n ↛ when n is an odd integer ≥ 3.

Page 414 line 1 — 20 Apr 1998

of t primes ↛ of t primes

▸**Page 416** line 11 of exercise 41 — 21 Jan 1998

$f(1000, 10)$ ↛ $f_2(1000, 10)$

▸**Page 416** line 3 of exercise 43 — 27 Apr 2001

$y \in S_m$ ↛ $y \in Q_m$

▸**Page 424** line 18 — 28 Oct 2000

Pablo Nuñez ↛ Pedro Nuñez

▸**Page 432** line 2 after (22) — 23 Jun 1998

$b_6^3 \cdot b_6^3 \cdot b_6^3$ ↛ $b_6^3 \cdot b_6^3$

Page 434 line 21 — 22 Aug 2002

(Paris: 1835) ↛ (Paris, 1835)

▸**Page 436** line 7 of exercise 17 — 03 Mar 2000

$\deg(u)$ ↛ $\deg(U)$

Page 438 exercise 20 — 29 Dec 2008

Investigate the accuracy ↜↝ Investigate *approximate* polynomial gcds and the accuracy

Page 441 in steps B1–B4 — 06 Jan 2011

[Summarize each step, as in the usual format for algorithms.]

▸**Page 442** line 14 — 06 Mar 1998

Pěstováni ↜↝ *Pěstování*

▸**Page 442** line −4 — 06 Jun 1999

3, ,. . . ↜↝ 3, . . .

Page 447 in steps D1–D3 (continuing to page 448) — 06 Jan 2011

[Summarize each step, as in the usual format for algorithms.]

Page 449 near the top — 20 Jul 2000

lines 5–8: A faster . . . arithmetic ↜↝ Faster algorithms for distinct-degree factorization when p is very large have been found by J. von zur Gathen, V. Shoup, and E. Kaltofen; the running time is $O(n^{2+\epsilon} + n^{1+\epsilon} \log p)$ arithmetic

lines 10–11: [See . . . to appear.] ↜↝ [See *Computational Complexity* **2** (1992), 187–224; *J. Symbolic Comp.* **20** (1995), 363–397; *Math. Comp.* **67** (1998), 1179–1197.]

Page 449 middle — 22 Aug 2002

line 15: *Paris* (1785) ↜↝ (Paris, 1785)
lines 24–25: *Sci. Paris*, series 2, **35** (1866) ↜↝ *Sci.*, series 2, **35** (Paris, 1866)
line 34: *Paris* **A291** (1980) ↜↝ **A291** (Paris, 1980)

Page 449 bottom line, and top of page 450 — 17 Jun 2002

extended by an astronomer . . . von Schubert's method independently ↜↝ extended by N. Bernoulli in 1708 and, more explicitly, by an astronomer named Friedrich von Schubert in 1793, who showed how to find all factors of degree n in a finite number of steps; see M. Mignotte and D. Ştefănescu, *Revue d'Hist. Math.* **7** (2001), 67–89. L. Kronecker rediscovered their approach independently,

▸**Page 451** line −13 — 28 Oct 2000

the range $(-\frac{169}{2}, \frac{169}{2})$ ↜↝ the range $(-\frac{169}{2} \,..\, \frac{169}{2})$

Page 452 in steps F1–F3 — 06 Jan 2011

[Summarize each step, as in the usual format for algorithms.]

▸**Page 455** lines 27 and 28 — 30 Sep 1998

Efficient Polynomial Computations ↜↝ *Effective Polynomial Computation*

▸**Page 457** line 3 of exercise 20 — 05 Feb 1998

$u(x) = (x - \alpha)u(x)$ and $v(x) = (\bar{\alpha}x - 1)u(x)$ ↜↝ $u(x) = (x - \alpha)w(x)$ and $v(x) = (\bar{\alpha}x - 1)w(x)$

▸**Page 458** line 11 — 14 Apr 1998

whenever $v_j \neq 0$. ↜↝ whenever $u_j \neq 0$.

▸**Page 458** line 20 — 06 Jun 1999

1^1 ↜↝ 1^2

▸**Page 459** line 3 of exercise 29 — 01 May 2001

$1/2 - 1/(2p^d)$ ↜↝ $1/2 - 1/(2p^{2d})$

▸**Page 461** line −18 — 21 Sep 2004

before 200 B.C. in Piṅgala's Hindu classic *Chandaḥsūtra* ↜↝ before A.D. 400 in Piṅgala's Hindu classic *Chandaḥśāstra*

[also change "next 1000 years" to "next several centuries", a few lines down]

Page 462 better wording in step A2 — 24 Jan 2003

Set N ... skip ↜↝ Set $t \leftarrow N \bmod 2$ and $N \leftarrow \lfloor N/2 \rfloor$. If $t = 0$, skip

Page 464 line −2 — 17 Feb 2001

$100,000$ ↜↝ 100,000

Page 465 line −6 — 07 Feb 2000

Our goal ↜↝ Thus $l(1) = 0$, $l(2) = 1$, $l(3) = l(4) = 2$, etc. Our goal

Page 466 line 3 — 02 Dec 2005

179 ↜↝ 179, 199

Page 466 lines 19–21 — 25 May 2008

Furthermore, as E. G. Thurber ... steps earlier. ↜↝ Furthermore, we can omit all the even numbers (except 2) in the first row, if we bring values of the form $d_j/2^e$ into the computation e steps earlier. [See E. Wattel and G. A. Jensen, *Math. Centrum Report* ZW1968-001 (1968), 18 pp.; E. G. Thurber, *Duke Math. J.* **40** (1973), 907–913.]

Page 468 line 9 — 06 Jan 2011

Corollary. ↜↝ **Corollary A.**

▸**Page 470** line 28 — 20 Sep 1998

$2^m + 2^{m+1} + 2^{d+1} + 2^d$ ↜↝ $2^m + 2^{m-1} + 2^{d+1} + 2^d$

Page 472 line −5 — 06 Jan 2011

Corollary. ↜↝ **Corollary E.**

Page 475 line 14 — 23 Oct 2006

the multisets M_{ij} and S_i, ↜↝ the multisets M_j, S_i, M_{ij}

Page 477 line 15 — 30 Jun 2002

des math. ↜↝ *des Math.*

Page 477 line 23 — 15 Sep 2005

948 cases ↜↝ 102 cases

Page 477 lines 26–28 or 26–29 — 12 Jun 2008

for all $k \geq 5$. ... Kevin R. Hebb ↯ for all $k \geq 5$. Neill Clift found in 2007 that $l(n) = l(2n) = l(4n) = 31$ when $n = 30958077$; and in 2008, astonishingly, he discovered that $l(n) > l(2n) = 34$ when $n = 375494703$. Kevin R. Hebb

Page 477 line 28 — 08 May 2007

Kevin ↯ Neill Clift found in 2007 that $l(n) = l(2n) = l(4n) = 31$ when $n = 30958077$. Kevin

Page 477 replacement for the bottom ten lines — 17 Nov 2008

r	$c(r)$	r	$c(r)$	r	$c(r)$
1	2	14	1087	27	2211837
2	3	15	1903	28	4169527
3	5	16	3583	29	7624319
4	7	17	6271	30	14143037
5	11	18	11231	31	25450463
6	19	19	18287	32	46444543
7	29	20	34303	33	89209343
8	47	21	65131	34	155691199
9	71	22	110591	35	298695487
10	127	23	196591	36	550040063
11	191	24	357887	37	994660991
12	379	25	685951	38	1886023151
13	607	26	1176431	39	3502562143

▸**Page 478** line 4 — 01 Feb 1999

[The ↯ The

Page 478 line 5 — 27 May 2008

Bleichenbacher. ↯ Bleichenbacher, and $c(29)$ through $c(33)$ by Neill Clift.

Page 478 replacement for lines 19–25 — 24 May 2008

table lists the first few values of $d(r)$, according to Flammenkamp and Clift:

r	$d(r)$	r	$d(r)$	r	$d(r)$	r	$d(r)$	r	$d(r)$	r	$d(r)$
1	1	6	15	11	246	16	4490	21	90371	26	1896704
2	2	7	26	12	432	17	8170	22	165432	27	3501029
3	3	8	44	13	772	18	14866	23	303475	28	6465774
4	5	9	78	14	1382	19	27128	24	558275	29	11947258
5	9	10	136	15	2481	20	49544	25	1028508	30	22087489

Page 478 the three lines following (49) — 13 May 2009

Computer calculations ... for $n = 32$. ↯ Computer calculations by Neill Clift show that $l(2^n - 1)$ is exactly equal to $n - 1 + l(n)$ for $1 \leq n \leq 64$; and E. G. Thurber [*Discrete Math.* **16** (1976), 279–289] has shown that several of these values, including the case $n = 32$, can actually be calculated by hand.

Page 479 line 4 after Table 1 — 27 Dec 2003

certain of the elements ↯ some of the elements

Page 479 lines −23 thru −20 — 17 Sep 2005

The chain ... theorem due to Hansen: ↝ Hansen pointed out that the chain constructed in Theorem F is an l^0-chain (see exercise 22); and he also established the following improvement of Eq. (50):

Page 479 new lines to be inserted at the bottom — 17 Sep 2005

Computations by Neill Clift have shown that $l(n) < l^0(n)$ when $n = 5784689$ (see exercise 42). This is the smallest case where Eq. (49) remains in doubt.

Page 481 line 1 of exercise 5 — 11 Jan 2003

$(k+1)$-st ↝ $(k+1)$st

Page 482 line 2 of exercise 14 — 30 Oct 2008

less than $m-1$. ↝ less than $m-1$, where $m = \lambda(a_{r-1})$.

▸**Page 483** line 9 of exercise 19 — 18 Dec 2000

$\prod_{p\in N} = n$ ↝ $\prod_{p\in N} p = n$

Page 483 exercise 27 — 15 Oct 2008

line 1: *M21* ↝ *M23*
line 2: six ↝ eight

Page 485 line −4 — 01 Jan 2002

105–107 ↝ 97–107

Page 485 replacement for exercise 42 — 17 Sep 2005

42. [*22*] (Neill Clift, 2005.) Show that neither 1, 2, 4, 8, 16, 32, 64, 65, 97, 128, 256, 353, 706, 1412, 2824, 5648, 11296, 22592, 45184, 90368, 180736, 361472, 361537, 723074, 1446148, 2892296, 5784592, 5784689 nor its dual is an l^0-chain.

43. [*M50*] Is $l(2^n-1) \le n-1+l(n)$ for all integers $n > 0$? Does equality always hold?

Page 489 line −14 — 07 Jun 1999

$n-1$ divisions ↝ $n-1$ divisions, since $v_n = u_n$

▸**Page 493** line 5 — 07 Jun 1999

$\lceil n/2 \rceil$ ↝ $\lceil n/2 \rceil - 1$

Page 501 line 3 after (38) — 13 Nov 1997

[remove the thin line above "F. Yates"]

Page 502 lines 8–18 — 23 Jun 2001

line 8: the *Walsh transform* ↝ the *Hadamard transform* or the *Walsh transform*
line 9: by J. L. Walsh ↝ by J. Hadamard [*Bull. Sci. Math.* (2) **17** (1893), 240–246] and by J. L. Walsh
lines 16–18: (See H. F. Harmuth, ... Walsh coefficients.) ↝ (See Section 7.2.1.1 for further discussion of the Hadamard–Walsh coefficients.)

▸**Page 504** replacement for bottom line — 28 Jun 1999

$$u_{[n]}(x) = \left(\frac{y_0 w_0}{x-x_0} + \cdots + \frac{y_n w_n}{x-x_n}\right) \Big/ \left(\frac{w_0}{x-x_0} + \cdots + \frac{w_n}{x-x_n}\right), \qquad (45)$$

▸**Page 507** replacement for bottom line, in 4th printing only — 10 Oct 1999

$$A=\begin{pmatrix}1&0&\bar1&0&0&1&\bar1\\0&1&0&0&0&1&0\\0&0&1&0&1&\bar1&1\\0&0&0&1&1&\bar1&1\end{pmatrix},\quad B=\begin{pmatrix}1&0&0&\bar1&\bar1&0&1\\0&1&0&1&0&0&0\\0&0&1&1&1&0&\bar1\\0&0&\bar1&\bar1&0&1&1\end{pmatrix},\quad C=\begin{pmatrix}1&1&0&0&0&0&0\\1&0&1&1&0&0&1\\1&0&0&0&1&1&1\\1&0&1&0&1&0&1\end{pmatrix}.\quad(53)$$

▸**Page 521** line 2 of exercise 56 — 26 Apr 2000

$\tau_{ijk}x_ix_k$ ↜ $\tau_{ijk}x_ix_j$

▸**Page 521** line 3 of exercise 60 — 26 Jan 2001

and only ↜ and only if

▸**Page 522** line 14 — 21 Apr 2002

problems of size $m\times n\times s$ ↜ problems of the respective sizes $m\times n\times s$ and $s\times m\times n$

▸**Page 523** line 15 of exercise 67 — 28 Jun 1999

))) ↜))

Page 526 in the line after (4) — 18 Jun 2010

nonnegative integer. From (4) ↜ nonnegative integer. If α itself is not an integer, there's more than one possibility for $V_m^\alpha z^{\alpha m}$ here.

From (4)

▸**Page 529** a spurious line just before Algorithm N — 07 Feb 2005

N ↜ (this error appeared in the 15th, 16th, and 17th printings only)

▸**Page 531** line 7 — 09 Aug 1999

$U=$ ↜ $U(z)=$

Page 536 line 1 of exercise 26 — 30 Jun 2002

$U_2z^2\cdots$ ↜ $U_2z^2+\cdots$

Page 539 replacement for the top three lines — 18 Jun 2008

It is almost legitimate to take ten readings $(m_0,\ldots,m_9)$ and then to output j if m_j is strictly less than m_i for all $i\neq j$; try again if the minimum value appears more than once. (See (h).) However, the parameter m isn't really constant in the real world.

Page 539 line 8 — 22 Nov 2006

horse. ↜ horse (unless you know, say, the jockey).

Page 539 line 10 of answer 6 — 08 Jun 2005

$f(x)=ax$.] ↜ $f(x)=ax$. See G. Frobenius, *Sitzungsberichte preußische Akademie der Wissenschaften* (1895), 82–83.]

Page 539 simplification for line −10 — 11 Jun 1998

$\lambda\delta_{\mu0}+\mu+\lambda-1-((\mu+\lambda-1)\bmod\lambda)$ ↜ $\lambda(\lceil\mu/\lambda\rceil+\delta_{\mu0})$

Page 540 line 12 — 25 Dec 2002

$e(n)=$ ↜ $e(n)=\rho(n+1)=$

Page 540 line −2 — 09 Nov 2002

that $\lambda = 1, \mu = 1$, or $\lambda = 2, \mu = 0$, is $\leadsto$ that $(\mu, \lambda) = (1, 1)$ or that $(\mu, \lambda) = (0, 2)$ is

Page 541 line 9 of answer 12 — 12 Jul 2003

its Applications $\leadsto$ *Its Applications*

Page 542 new answer — 08 May 2005

19. Clearly $\Pr(\text{no final cycle has length } 1) = (m-1)^m/m^m$. R. Pemantle [*J. Algorithms* **54** (2005), 72–84] has shown that $\Pr(\lambda = 1) = \Theta(m^{k/2})$, and that $\Pr((\mu+\lambda)^2 > 2m^k x$ and $\lambda/(\mu+\lambda) \le y)$ rapidly approaches ye^{-x}, when $x > 0$, $0 < y < 1$, and $m \to \infty$. The k-dimensional analogs of exercises 13 and 14 remain unsolved.

Page 542 update on answer 21 — 08 Mar 1998

Scott Fluhrer has discovered *another* fixed point of Algorithm K, namely the value 5008502835(!). He also found the 3-cycle

$$0225923640 \to 2811514413 \to 0590051662 \to 0225923640,$$

making a total of seven cycles in all. Exactly 128 starting numbers lead to the repeating value 5008502835; the smallest of these is 0008502835, the largest is 9944390948.

Page 543 line 2 of answer 5 — 24 Jan 2003

mod m, and in general, $\leadsto$ mod m; and in general, if $b = a - 1$,

▸**Page 546** line 3 — 03 Jul 2010

$9999999000^n \leadsto 999999900^n$

▸**Page 546** line 20 — 28 Sep 2002

word size $w \leadsto$ word size

▸**Page 546** line 24 — 24 Nov 2007

$+ b_n \leadsto - b_n$

▸**Page 546** replacement for line −4 — 24 Nov 2007

$$x_n = (x_{n-3} - x_{n-10} - b_n) \bmod 10 = x_{n-3} - x_{n-10} - b_n + 10b_{n+1}$$

▸**Page 546** line −2 — 19 Feb 1999

$\bar{x}_{n-1}\bar{x}_n \leadsto \bar{x}_{n+1}\bar{x}_n$

▸**Page 547** line 8 of answer 14 — 01 Dec 2007

$a_j(x_{n-1}\ldots x_{n-k})_b \leadsto a_j(x_{n-1}\ldots x_{n-j})_b$

▸**Page 549** lines 3–6 — 08 Nov 2005

Now since ... so $\lambda = \lambda_1\lambda_2$. $\leadsto$ Now let λ' be the order of a_1a_2. Since $(a_1a_2)^{\lambda'} \equiv 1$, we have $1 \equiv (a_1a_2)^{\lambda'\lambda_1} \equiv a_2^{\lambda'\lambda_1}$; hence $\lambda'\lambda_1$ is a multiple of λ_2. This implies that λ' is a multiple of λ_2, since λ_1 is relatively prime to λ_2. Similarly, λ' is a multiple of λ_1; hence λ' is a multiple of $\lambda_1\lambda_2$. But obviously $(a_1a_2)^{\lambda_1\lambda_2} \equiv 1$, so $\lambda' = \lambda_1\lambda_2$.

Page 549 line 6 of answer 16 — 28 Mar 2005

distinct roots. $\leadsto$ distinct roots. [J. L. Lagrange, *Mém. Acad. Roy. Sci. Berlin* **24** (1768), 181–250, §10.]

Page 550 line 18 — 03 Jun 2001

is unfeasible $\leadsto$ isn't feasible

▸**Page 550** line 19 — 13 Jan 1998

succeed in find $\leadsto$ succeed in finding

Page 551 replacement for line 2 — 27 Feb 2004

$$Y_{n+2} \approx (10 + 6(Y_{n+1} + \epsilon_1) - 9(Y_n + \epsilon_2)) \mod 20,$$

▸**Page 553** line -2 — 18 Feb 2004

$X_{n+p^{f+1}} \leadsto X_{n+p^{f+1}d}$

▸**Page 555** line 10 — 01 Jun 2001

$a_1X_{(n-1)1} + a_2X_{(n-2)2} + X_{(n-2)3} \leadsto a_1X_{(n-1)1} + a_2X_{(n-2)1} + X_{(n-2)3}$

▸**Page 555** line 1 of answer 18 — 03 Jun 2001

The third-most-significant $\leadsto$ By exercise 16, the most significant

Page 556 line 14 — 08 Jul 2001

$\langle sX_n + Y_n\rangle \leadsto \langle (X_n + Y_n) \bmod rs\rangle$

▸**Page 557** lines 3 and 4 — 25 Jan 2003

$i \leadsto j$ (three changes)

Page 557 last lines of answer 29 — 29 Jul 2006

S. Xie, ... 157–177 $\leadsto$ H. Fredricksen and J. Maiorana, *Discrete Math.* **23** (1978), 207–210, who essentially discovered Algorithm 7.2.1.1F.

▸**Page 557** line 1 of answer 30 — 28 Sep 2002

By exercise $\leadsto$ (a) By exercise

▸**Page 558** bottom line — 03 Feb 2005

$x - u_2 \leadsto x + u_2$ and $x - u_d \leadsto x + u_d$

▸**Page 565** line 5 of answer 24 — 05 Feb 2005

strings α, $\leadsto$ strings α, and we also have $\sum_{|\alpha|=t} N(\alpha) = n$;

▸**Page 566** last two lines of answer 24 — 03 Feb 2005

$K(i : (j : \alpha), (\beta : l) : i) - K(i-1 : (j : \alpha), (\beta : l+1) : i-1)$, where $i = \min(t+k, t-l)$ and $j = \max(0, k+l)$. $\leadsto$ $K(i : (\alpha : i-k), i : (\beta : i+l)) - K(i-1 : (\alpha : i-k), i-1 : (\beta : i+l))$, where $i = \min(t+k, t-l)$.

▸**Page 566** lines -9, -8, and -7 — 18 Sep 2002

Whitworth, who stated the question as problem 667 in *Choice and Chance* (Cambridge, 1867) and gave the solution in $\leadsto$ Whitworth, in problem 667 of

▸**Page 569** line 10 — 12 Apr 1999

$-ic^3t^3 \leadsto -ic_3t^3$

▸**Page 569** line 12 — 07 Jan 2004

for $j \geq 8$ $\leadsto$ for $j \geq 3$

▸**Page 570** line 1 — 12 Apr 1999

(1093) ↜↝ (1953)

Page 571 improvement to answer 32 — 29 Mar 2005

Not if they ... equal strength ↜↝ Not if, say, X and Y each take the values $(-n, m)$ with the respective probabilities $(m/(m+n), n/(m+n))$, where $m < n < (1+\sqrt{2})m$. [Suppose two competitors differ by X after playing one round of golf. Then they are of equal strength

Page 572 new answer — 13 Jun 2001

[Renumber answer 3 as answer 2, delete the former answer 2, and insert the following new material:]

3. The sum is $((2^n x)) - ((x))$. [See *Trans. Amer. Math. Soc.* **65** (1949), 401.]

Page 573 line 3 of answer 9 — 28 Sep 2002

math. ↜↝ *Math.*

▸**Page 577** replacement for last three lines of answer 3 — 04 Jul 2003

$$m = 9,\quad a = 4 \text{ or } 7,\quad \nu_2^2 = \nu_3^2 = 5;$$
$$m = 9q,\quad a = 3q+1 \text{ or } 6q+1,\quad \nu_4^2 = 2.$$

▸**Page 578** line 4 of answer 8 — 16 Feb 1999

the bound (39) ↜↝ the bound (40)

▸**Page 581** in answer 20 — 25 Jul 2005

lines 2 and 3: plus $(X_0 \bmod 4)/2^e$, where $\sigma(y) = (ay + \lfloor a/4 \rfloor) \bmod 2^{e-2}$. ↜↝
plus $2^{-e}(t, \ldots, t)$, where $\sigma(y) = (ay + \lfloor a/4 \rfloor t) \bmod 2^{e-2}$ and $t = X_0 \bmod 4$.
line 9: $(-ay - \lceil a/4 \rceil) \bmod 2^{e-2}$ ↜↝ $(-ay - \lceil a/4 \rceil t) \bmod 2^{e-2}$

Page 582 new sentence to append to answer 26 — 06 Feb 2005

Since $m/d > 1$, we can also replace N by $N \bmod (m/d)$. [And *delete* the first sentence, 'When $m = 1$, ... zero.']

▸**Page 584** line 4 of answer 3 — 20 Aug 2001

$U > k\lfloor m/k \rfloor$ ↜↝ $U \geq k\lfloor m/k \rfloor$

Page 585 improvement to Fig. A–4 — 04 Dec 2005



▸**Page 585** line −3 — 04 Oct 1999

absolute minimum ↜↝ absolute maximum

Page 587 in answer 19 — 06 Jan 2011

B1 $\nwarrow\!\!\!\!\searrow$ **N1** and **B2** $\nwarrow\!\!\!\!\searrow$ **N2** (to avoid conflict with answer 27)

Page 587 replacement for lines −13 and −12 — 29 Jun 1999

N2. [Iterate.] If $U \geq r$, set $N \leftarrow N+1$, $q \leftarrow q(1-p)(t-1+N)/N$, $r \leftarrow r+q$, and repeat this step. Otherwise return N and terminate. ∎

▸**Page 587** bottom line — 03 Feb 2005

$\frac{2}{5}(a-bu) - \frac{2}{3}$ $\nwarrow\!\!\!\!\searrow$ $\frac{2}{5}(a-bu) - \frac{2}{3}a$

▸**Page 588** line 1 — 03 Feb 2005

$I = \int_0^{a/b} u\sqrt{a-bu}\,du = \frac{4}{15}a^{5/2}/b^2$ $\nwarrow\!\!\!\!\searrow$ $I = 2\int_0^{a/b} u\sqrt{a-bu}\,du = \frac{8}{15}a^{5/2}/b^2$

▸**Page 588** line −2 of answer 21 — 03 Feb 2005

$E = \int_0^{r(x)}$ $\nwarrow\!\!\!\!\searrow$ $E = 2\int_0^{r(x)}$

Page 588 last line of answer 22 — 29 Jun 1999

208]. $\nwarrow\!\!\!\!\searrow$ 208.]

Page 588 beginning of answer 25 — 19 May 2005

line 1: ∨ $\nwarrow\!\!\!\!\searrow$ |
line 2: ∧ $\nwarrow\!\!\!\!\searrow$ &

▸**Page 589** line 3 of answer 29 — 25 Mar 2006

David Seneschol $\nwarrow\!\!\!\!\searrow$ David Seneschal

Page 590 end of answer 31 — 08 Jul 1999

Preprint ... 1997) $\nwarrow\!\!\!\!\searrow$ *Lecture Notes in Comp. Sci.* **1470** (1998), 1–20

▸**Page 591** in printings 5–14 only — 07 Feb 2004

[the top line of page 591 was a duplicate of the bottom line of page 590]

▸**Page 591** lines −3 and −2 of answer 8 — 29 Jun 1999

decrease n and N by 1 $\nwarrow\!\!\!\!\searrow$ set $n \leftarrow n-1$, $N \leftarrow N-X-1$,

▸**Page 592** line 2 of answer 13 — 01 Jan 2008

$(x+1)$ $\nwarrow\!\!\!\!\searrow$ $(x-1)$

Page 592 new answer — 22 Nov 2003

18. (a) Oriented trees that essentially merge $(1, 2, \dots)$ with $(n, n-1, \dots)$, such as



(b) Collections of 1-cycles and 2-cycles. (c) Binary search trees on the keys $(1, 2, \dots, n)$, with k_j the parent of j (or j, at the root); see Section 6.2.2. The number of $(k_1, \dots, k_n)$ in each case is (a) 2^{n-1}; (b) $t_n \geq \sqrt{n!}$, see 5.1.4–(40); (c) $\binom{2n}{n}\frac{1}{n+1}$. [Case (a) represents the least common permutation; case (b) represents the most common, when $n \geq 18$. See D. P. Robbins and E. D. Bolker, *Æquationes Mathematicæ* **22** (1981), 268–292; D. Goldstein and D. Moews, *Æquationes Mathematicæ* **65** (2003), 3–30.]

Page 593 another new answer — 12 Jan 2009

19. See N. Duffield, C. Lund, and M. Thorup, *JACM* **54** (2007), 32:1–32:37.

▸**Page 593** line 1 of answer 6 — 19 Mar 2005

By exercise 5 ↜ By exercise 4

Page 596 line −5 — 29 Jun 1999

. We ↜ We

Page 599 lines 3 and 4 of answer 7 — 10 May 2009

related problems. See also L. Blum ... 364–383; J. Boyar ↜ related problems; see also J. Boyar

▸**Page 599** line 1 of answer 9 — 26 Jan 2001

The values ↜ (a) The values

Page 601 replacement for lines 15–41 — 01 Jan 2002

```
1     CONTINUE
      X(2)=X(2)+1
      SS=SSEED
      T=TT-1
10    DO 12 J=KK,2,-1
         X(J+J-1)=X(J)
12       X(J+J-2)=0
      DO 14 J=KKK,KK+1,-1
         X(J-(KK-LL))=X(J-(KK-LL))-X(J)
         IF (X(J-(KK-LL)) .LT. 0) X(J-(KK-LL))=X(J-(KK-LL))+MM
         X(J-KK)=X(J-KK)-X(J)
         IF (X(J-KK) .LT. 0) X(J-KK)=X(J-KK)+MM
14    CONTINUE
      IF (MOD(SS,2) .EQ. 1) THEN
         DO 16 J=KK,1,-1
16          X(J+1)=X(J)
         X(1)=X(KK+1)
         X(LL+1)=X(LL+1)-X(KK+1)
         IF (X(LL+1) .LT. 0) X(LL+1)=X(LL+1)+MM
       END IF
```

Page 602 (now page 601) two new lines before 'END' — 01 Jan 2002

```
        DO 22 J=1,10
22         CALL RNARRY(X,KKK)
```

Page 602 line 8 of answer 11 — 01 Jan 2002

copy in *ul* ↜ copy in x

Page 602 from line 28 of answer 11 to that answer's end — 01 Jan 2002

```
double ss=2.0*ulp*((seed&0x3fffffff)+2);
for (j=0;j<KK;j++) {
  x[j]=0; u[j]=ss;                        /* bootstrap the buffer */
  ss+=ss; if (ss>=1.0) ss-=1.0-2*ulp;  /* cyclic shift of 51 bits */
```

```
  }
  x[1]=1; u[1]+=ulp;              /* make u[1] (and only u[1]) "odd" */
  for (s=seed&0x3fffffff,t=TT-1; t; ) {
    for (j=KK-1;j>0;j--)
      x[j+j]=x[j],u[j+j]=u[j],x[j+j-1]=0,u[j+j-1]=0.0;  /* "square" */
    for (j=KK+KK-2;j>=KK;j--) {
      x[j-(KK-LL)]=x[j]^x[j-(KK-LL)],
        u[j-(KK-LL)]=mod_sum(u[j-(KK-LL)],u[j]);
      x[j-KK]=x[j]^x[j-KK],u[j-KK]=mod_sum(u[j-KK],u[j]);
    }
    if (is_odd(s)) {                              /* "multiply by z" */
      for (j=KK;j>0;j--)  x[j]=x[j-1],u[j]=u[j-1];
      x[0]=x[KK],u[0]=u[KK];          /* shift the buffer cyclically */
      x[LL]=x[KK]^x[LL],u[LL]=mod_sum(u[LL],u[KK]);
    }
    if (s) s>>=1; else t--;
  }
  for (j=0;j<LL;j++) ran_u[j+KK-LL]=u[j];
  for (;j<KK;j++) ran_u[j-LL]=u[j];
  for (j=0;j<10;j++) ranf_array(u,KK+KK-1);   /* warm everything up */
}
int main() {
  register int m; double a[2009];             /* a rudimentary test */
  ranf_start(310952);
  for (m=0;m<2009;m++) ranf_array(a,1009);
  printf("%.20f\n", ran_u[0]);            /* 0.36410514377569680455 */
  ranf_start(310952);
  for (m=0;m<1009;m++) ranf_array(a,2009);
  printf("%.20f\n", ran_u[0]);            /* 0.36410514377569680455 */
  return 0;
}
```

Page 603 last line of answer 12 — 26 Dec 2002

9387 ↝ 24130

Page 604 fourth-from-last line of the program in answer 15 — 29 Mar 2005

`ran_arr_buf[100]` ↝ `ran_arr_buf[KK]`

Page 604 insert new line at end of answer 15 — 01 Jan 2002

Reset *ran_arr_ptr* = &*ran_arr_sentinel* if *ran_start* is used again.

▸**Page 605** last line of answer 10 — 15 Feb 2006

any infinite sequence of integers with $k_{j+1} > k_j$ ↝
any doubly infinite sequence of integers with $k_{j+1} > k_j$ and $k_0 = 0$.

Page 605 line 2 of answer 12 — 26 May 2008

exercise 7.1–18 ↝ exercise 7.1.3–7

Page 606 line 3 of answer 18 — 09 Jun 2003

$y_1, y_2, \ldots$ ↝ $\{y_1, y_2, \ldots\}$

▸**Page 606** line 13 of answer 18 — 09 Jun 2003

$|x|$ and $|y|$ are fixed $\leadsto$ $|x|$ and $|y|$ are sufficiently small

▸**Page 607** line 2 of answer 21 — 26 Aug 1999

the test $\leadsto$ the text

▸**Page 607** line 3 of answer 21 — 26 Jan 2007

the systems $\leadsto$ the system

▸**Page 608** last line of answer 24 — 26 Jan 2001

digits.] $\leadsto$ digits.

Page 608 line −2 of answer 25 — 26 Jan 2007

aaaaa $\leadsto$ *a*

Page 608 bottom line — 03 Jun 2005

induction. $\leadsto$ induction. [A. D. Booth, in *Quarterly J. Mechanics and Applied Math.* **4** (1951), 236–240, applied this principle to two's complement multiplication.]

▸**Page 610** last two lines of answer 31(c) — 26 Jan 2007

$2^{-\mu}u$ is an integer. $\leadsto$ $2^r u$ is an integer, for sufficiently large r.

Page 611 line 3 of answer 34 — 15 Dec 2009

converted to α_n $\leadsto$ converted to this "canonical signed bit representation"

Page 611 additions to first paragraph of answer 34 — 15 Dec 2009

(See exercise 7.1.3–35.)

Page 611 replacement for answer 1 — 28 Jan 1999

1. $N = (62, +.60\ 22\ 14\ 00)$; $h = (37, +.66\ 26\ 10\ 00)$. Note that the quantity $10h$ would be $(38, +.06\ 62\ 61\ 00)$.

▸**Page 611** line 3 of answer 5 — 02 May 2000

$x \sim y \leadsto bx \sim by$

▸**Page 612** line 2 of answer 12 — 28 Mar 2006

we have $1/b <$ $\leadsto$ we have $1/b \leq$

▸**Page 613** replacement for answer 16 — 04 Oct 2010

16. If $|c| \geq |d|$, then set $r \leftarrow d \oslash c$, $s \leftarrow c \oplus (r \otimes d)$; $x \leftarrow \big(a \oplus (b \otimes r)\big) \oslash s$, $y \leftarrow \big(b \ominus (a \otimes r)\big) \oslash s$. Otherwise set $r \leftarrow c \oslash d$, $s \leftarrow d \oplus (r \otimes c)$; $x \leftarrow \big((a \otimes r) \oplus b\big) \oslash s$, $y \leftarrow \big((b \otimes r) \ominus a\big) \oslash s$. Then $x + iy$ is the desired approximation to $(a + bi)/(c + di)$. Computing $s' \leftarrow 1 \oslash s$ and multiplying twice by s' may be better than dividing twice by s. As with (11), gradual underflow is recommended for the calculation of r unless special precautions are taken. [*CACM* **5** (1962), 435. Other algorithms for complex arithmetic and function evaluation are given by P. Wynn, *BIT* **2** (1962), 232–255. For $|a + bi|$, see Paul Friedland, *CACM* **10** (1967), 665.]

▸**Page 613** replacement for answer 19 — 10 Nov 2002

19. $(73-(5-[\text{rounding digits are } \frac{b}{2}0\ldots0])(6-[\text{magnitude is rounded up}])+[e_v<e_u]+[\text{first rounding digit is } \frac{b}{2}]-[\text{fraction overflow}]-10[\text{result zero}]+7[\text{rounding overflow}]+7N+(3+(16+[\text{result negative}])[\text{opposite signs}])X)u$, where N is the number of left shifts during normalization, and X is the condition that rX receives nonzero digits and there is no fraction overflow. The maximum time of $84u$ occurs for example when

$$u = -50\ 01\ 00\ 00\ 00, \quad v = +45\ 49\ 99\ 99\ 99, \quad b = 100.$$

[The average time, considering the data in Section 4.2.4, will be less than $47u$.]

▸**Page 614** line 3 of answer 14 — 01 Mar 2007

$b^{2-p} \mathrel{\not\rightarrow} (1+b^2)b^{-p}$

▸**Page 615** line 1 of answer 16 — 01 Mar 2007

101.11111 $\mathrel{\not\rightarrow}$ 101.11110

Page 615 line -1 — 27 Jun 2003

For another $\mathrel{\not\rightarrow}$ Kahan observed that $s_n \ominus c_n = \sum_{k=1}^{n}(1+\phi_k)x_k$ where $|\phi_k| \le 2\epsilon + O((n+1-k)\epsilon^2)$. For another

Page 616 line 1 — 27 Jan 2000

can combine them $\mathrel{\not\rightarrow}$ may be able to match them up

▸**Page 616** line 5 of answer 21 — 09 Sep 1999

$e_u = e_u = p \mathrel{\not\rightarrow} e_u = e_v = p$

▸**Page 617** line 1 of answer 27 — 31 Jul 1998

$1 \oslash u) \oslash u \mathrel{\not\rightarrow} 1 \oslash (1 \oslash u)$

▸**Page 620** near the top — 25 Feb 2010

line 3: $\sum_{m\le 0} \mathrel{\not\rightarrow} \sum_{m<0}$
line 4: $F(10^k 4^k) \mathrel{\not\rightarrow} F(4^k)$

Page 622 line 14 of answer 18 — 22 Mar 2002

$p(b^{n-1}a, b^n) \mathrel{\not\rightarrow} p(b^{n-1}a, b^n)$

▸**Page 622** answer 19 — 27 Sep 1999

line 1: $\log 10F_n \mathrel{\not\rightarrow} \log_{10} F_n$
line 2: $\log 10\phi \mathrel{\not\rightarrow} \log_{10} \phi$

Page 625 in answer 16 — 30 Jan 2010

[change the step numbers S1, S2, S3 respectively to Q1, Q2, Q3]

Page 625 last line of answer 22 — 07 Apr 2003

$b = 10.) \mathrel{\not\rightarrow} b = 10$. Similarly, when $b = 2^{16}$ we can let $u = (\texttt{7fff800100000000})_{16}$, $v = (\texttt{800080020005})_{16}$.)

▸**Page 625** replacement for answer 23 — 23 Jan 1998

23. Obviously $v\lfloor b/(v+1)\rfloor < (v+1)\lfloor b/(v+1)\rfloor \le b$; and the lower bound certainly holds if $v \ge b/2$. Otherwise $v\lfloor b/(v+1)\rfloor \ge v(b-v)/(v+1) \ge (b-1)/2 > \lfloor b/2\rfloor - 1$.

▸**Page 626** replacement for lines 1–5 of answer 25 — 23 Jan 1998

25.
```
002      ENTA 1           1
003      ADD  V+N-1       1
004      STA  TEMP        1
005      ENTA 1           1
006      JOV  1F          1   Jump if v_{n-1} = b - 1.
007      ENTX 0           1
008      DIV  V+N-1       1   Otherwise compute ⌊b/(v + 1)⌋.
009      JOV  DIVBYZERO   1   Jump if v_{n-1} = 0.
010  1H  STA  D           1
```
[All subsequent line numbers for Program D should now be increased by 4.]

▸**Page 627** line 1 — 25 Feb 1998

convex function ↜ concave function

Page 629 last lines of answer 40 — 28 Dec 2000

[See T. Jebelean, ... 459.] ↜ [See A. Schönhage and E. Vetter, *Lecture Notes in Comp. Sci.* **855** (1994), 448–459; W. Krandick and T. Jebelean, *J. Symbolic Computation* **21** (1996), 441–455.]

Page 629 line 5 of answer 41 — 02 May 2006

trial divisor.) ↜ trial divisor. To compute w' when b is a power of 2, notice that if $w_0w' \equiv 1$ (modulo 2^e) then $w_0w'' \equiv 1$ (modulo 2^{2e}) when $w'' = (2 - w_0w')w'$, by the 2-adic analog of "Newton's method.")

Page 629 line 10 of answer 41 — 10 Aug 1998

signed-magnitude ↜ signed magnitude

▸**Page 630** answer 43 — 08 Nov 2005

$xy/255$ ↜ $uv/255$ (twice)

▸**Page 631** displayed formula in answer 5 — 31 May 1999

p prime ↜ p prime

Page 631 new sentence at end of answer 5 — 05 Dec 2000

Replacing 100 by 256 and allowing even moduli gives $2^8 3^5 5^3 \ldots 251^1 \approx 1.67 \cdot 10^{109}$.

Page 632 last line of answer 14 — 07 Apr 2003

324.] ↜ 324. For improved error bounds, and extensions to moduli of the form $k \cdot 2^n \pm 1$, see Colin Percival, *Math. Comp.* **72** (2002), 387–395.]

▸**Page 632** line 4 of answer 1 — 11 Nov 1999

[shift '+03' one column to the left]

Page 635 line 2 of answer 1 — 08 Apr 2002

B_j system ↜ B_J system

▸**Page 636** replacement for answer 9 — 14 May 2006

9. Let $p_k = 2^{2^{k+2}}$. By induction on k we have $v_k(u) \le \frac{16}{5}(1 - 1/p_k)(\lfloor u/2 \rfloor + 1)$; hence $\lfloor v_k(u)/16 \rfloor \le \lfloor \lfloor u/2 \rfloor /5 \rfloor = \lfloor u/10 \rfloor$ for all integers $u \ge 0$. Furthermore, since

$v_k(u+1) \ge v_k(u)$, the smallest counterexample to $\lfloor v_k(u)/16 \rfloor = \lfloor u/10 \rfloor$ must occur when u is a multiple of 10.

Now let $u = 10m$ be fixed, and suppose $v_k(u) \bmod p_k = r_k$ so that $v_{k+1}(u) = v_k(u) + (v_k(u) - r_k)/p_k$. The fact that $p_k^2 = p_{k+1}$ implies that there exist integers m_0, m_1, m_2, ... such that $m_0 = m$, $v_k(u) = (p_k - 1)\,m_k + x_k$, and $m_k = m_{k+1}p_k + x_k - r_k$, where $x_{k+1} = (p_k + 1)x_k - p_k r_k$. Unwinding this recurrence yields

$$v_k(u) = (p_k - 1)\,m_k + c_k - \sum_{j=0}^{k-1} p_j r_j \prod_{i=j+1}^{k-1} (p_i + 1), \qquad c_k = 3\,\frac{p_k - 1}{p_0 - 1}.$$

Furthermore $v_k(u) + m_k = v_{k+1}(u) + m_{k+1}$ is independent of k, and it follows that $v_k(u)/16 = m + (3 - m_k)/16$. So the minimal counterexample $u = 10y_k$ is obtained for $0 \le k \le 4$ by setting $m_k = 4$ and $r_j = p_j - 1$ in the formula $y_k = \frac{1}{16}(v_k + m_k - c_0)$. In hexadecimal notation, y_k turns out to be the final 2^k digits of `434243414342434`.

Since $v_4(10y_4)$ is less than 2^{64}, the same counterexample is also minimal for all $k > 4$. One way to work with larger operands is to modify the method by starting with $v_0(u) = 6\lfloor u/2 \rfloor + 6$ and letting $c_k = 6(p_k - 1)/(p_0 - 1)$, $m_0 = 2m$. (In effect, we are truncating one bit further to the right than before.) Then $\lfloor v_k(u)/32 \rfloor = \lfloor u/10 \rfloor$ when u is less than $10z_k$, for $1 \le k \le 7$, where $z_k = \frac{1}{32}(v_k + m_k - 6)$ when $m_k = 7$, $r_0 = 14$, and $r_j = p_j - 1$ for $j > 0$. For example, $z_4 =$ `1c342c3424342c34`. [This exercise is based on ideas of R. A. Vowels, *Australian Comp. J.* **24** (1992), 81–85.]

Page 638 last two lines of answer 18 — 15 Apr 1999

(Is this ... 10^4?) $\nwarrow$ (The *smallest* example is actually round$((.1111111001)_2 \cdot 2^{784}) = .1011 \cdot 10^{236}$, round$(.1011 \cdot 10^{235}) = (.11111110010)_2 \cdot 2^{784}$, found by Fred J. Tydeman.)

Page 638 answer 4.5.1–1 — 15 Jan 2002

positive. $\nwarrow$ positive. (See also the answer to exercise 4.5.3–39.)

▸**Page 641** last line of answer 10 — 18 Feb 2004

for arbitrary f and g, when $m \le n$. $\nwarrow$ when $m \le n$, $f(m) = O(m)$, and $g(n) = O(n)$.

▸**Page 641** first line of answer 14 — 18 Feb 2004

$(2^p - 1)$ $\nwarrow$ $(p^2 - 1)$

▸**Page 642** line 2 of answer 17 — 29 Sep 2001

$u' = 1$ $\nwarrow$ $u' = u$

Page 645 new answer — 23 May 2008

32. Yes: See G. Maze, *J. Discrete Algorithms* **5** (2007), 176–186.

Page 645 new answer — 30 Jan 1998

34. Brigitte Vallée [*Algorithmica* **22** (1998), 660–685] has found an elegant and rigorous analysis of Algorithm B, using an approach quite different from that of Brent. Indeed, her methods are sufficiently different that they are not yet known to predict the same behavior as Brent's heuristic model. Thus the problem of analyzing the binary gcd algorithm, now solved rigorously for the first time, continues to lead to ever more tantalizing questions of higher mathematics.

Page 645 lines 4 and 5 of answer 36 — 03 Aug 2005

$u = \max(a_{n+2}, 2a_{n+1})$ and $v = a_{n+3} - u$ $\nwarrow$ $u = a_{n+2}$ and $v = u + (-1)^n$

▸**Page 646** in answer 39 — 29 Jun 1998

line 1 of step Y6: if $t_1 < 0$ ↝ if $t_1 \le 0$
line 2 after step Y6: $-u$ after ↝ $-u$, $0 < u_3 \le u$, $0 < v_3 \le v$ after

Page 646 in lines −8 and −4 of answer 39 — 01 May 2006

after step Y2 ↝ after step Y1 (twice)

▸**Page 647** line 1 of answer 7 — 06 Jan 2000

$F_k\, F_{n-k}$ ↝ $F_k\, F_{n+1-k}$

Page 649 line 8 — 17 Sep 1998

59–62. ↝ 59–62. Hurwitz also explained how to multiply by an arbitrary positive integer, in *Vierteljahrsschrift der Naturforschenden Gesellschaft in Zürich* **41** (1896), Jubelband II, 34–64, §2.]

Page 649 new sentence to conclude answer 15 — 18 Dec 2008

For further developments, see J. Vuillemin, *ACM Conf. LISP and Functional Programming* **5** (1988), 14–27.

▸**Page 650** line 20 — 12 Jan 2006

$n!\, z^n\, {}_2F_0(n+1, -n\,;;-1/z)$ ↝ $z^n\, {}_2F_0(n+1, -n\,;;-1/z)$

▸**Page 650** line −3 — 12 Jan 2006

$$\sum_{m\ge 0} z^m \;\leadsto\; \sum_{m\ge 0} \frac{z^m}{m!}$$

Page 652 replacement for answer 29 — 13 Jan 2000

29. We have $\sum_{n=1}^N n \ln n = \frac{1}{2}N^2 \ln N + O(N^2)$ by Euler's summation formula (see exercise 1.2.11.2–7). Also $\sum_{n=1}^N n \sum_{d\backslash n} \Lambda(d)/d = \sum_{d=1}^N \Lambda(d) \sum_{1\le k\le N/d} k$, and this is $O(\sum_{d=1}^N \Lambda(d) N^2/d^2) = O(N^2)$. Indeed, $\sum_{d\ge 1} \Lambda(d)/d^2 = -\zeta'(2)/\zeta(2)$.

Page 653 addition to end of answer 32 — 19 Mar 2007

paper.) ↝ paper. Incidentally, "Morse code" was really invented by F. C. Gerke in 1848; Morse's prototypes were quite different.)

▸**Page 653** corrections to answer 34 — 18 Feb 2004

line 5: $s(y) = \sum_{d\backslash y} k(y/d)$ ↝ $s(y) = \sum_{d\backslash y} \mu(d) k(y/d)$
line 6: $\sum_{y=1}^n [d\backslash y]$ ↝ $\sum_{y=1}^n \sum_{d\backslash y}$

Page 653 line 2 of answer 35 — 12 Jul 2003

its Applications ↝ *Its Applications*

▸**Page 654** line 1 of answer 39 — 08 Jan 2000

and no fraction with denominator < 287 lies in $//\,2, 1, 95\,//$ ↝ $//2, 1, 95//$

▸**Page 655** line 8 — 25 Jul 2000

6 (1860) ↝ **3** (1861)

Page 655 line −10 — 21 Jul 2009

$d_{4n+3} = 1$; the dragon curve ↝ $d_{4n+3} = 1$; the Jacobi symbol $(\frac{-1}{n})$ is $1 - 2d_n$. The dragon curve

▸**Page 656** last line of answer 45 — 18 May 2002

Theorem L ↜↝ Corollary L

▸**Page 657** line 15 — 31 May 1999

$(-1)^(j-1)$ ↜↝ $(-1)^{(j-1)}$

▸**Page 658** near the top — 14 May 1998

line 13: Henri Cohen has noticed ↜↝ Brent noticed
line 18: *A Course* ↜↝ See the analysis by Henri Cohen in *A Course*

▸**Page 658** answer 8 — 21 Aug 2001

line 2: $X[M]$ ↜↝ $X[M-1]$
lines 3 and 6: $k \le M$ ↜↝ $k < M$
line 7: $j \le M$ ↜↝ $j < M$

▸**Page 658** line 3 of answer 8 — 23 Apr 2001

$j \leftarrow 1$, ↜↝ $j \leftarrow 1$, $k \leftarrow 1$,

Page 659 line 9 — 18 Oct 2006

Section 7.1 ↜↝ Section 7.1.3

▸**Page 659** answer 9 — 18 May 2002

line 8: less than N [see ↜↝ less than N; see
line 9: 722]. ↜↝ 722.]

▸**Page 661** line 3 of answer 17 — 20 Apr 2009

SICOMP **7** (1975) ↜↝ *SICOMP* **4** (1975)

Page 661 lines 3 and 4 of answer 22 — 03 May 2002

$p_n = (b_n - 1)/(n-2)$ ... not prime. ↜↝ $p_n = [n \text{ nonprime}](b_n - 1)/(n-2) < [n \text{ nonprime}]b_n/(n-1)$, where b_n is the number of bad x such that $1 \le x < n$.

▸**Page 662** last line of answer 22 — 06 Jun 1999

remarks.] ↜↝ remarks.

Page 663 line 5 of answer 27 — 22 Aug 2002

Paris **87** (1878) ↜↝ **87** (Paris, 1878)

Page 663 line 9 of answer 27 — 07 Apr 2003

a single-precision number. ↜↝ small. (See also exercise 4.3.2–14.)

Page 664 line 6 — 23 Nov 1998

Comp., to appear. ↜↝ *Comp.* **67** (1998), 1735–1738.

▸**Page 664** line 1 of answer 29 — 26 Jan 2001

solutions to the equation ↜↝ solutions to the inequality

▸**Page 664** line 4 of answer 32 — 18 Feb 2004

$M^2 - M \le y \le M^2$ ↜↝ $M^2 - M \le y < M^2$

▸**Page 664** line −2 — 18 Feb 2004

$(\frac{\bar{x}}{N}) = 1$ ↜↝ $(\frac{\bar{x}}{N}) \ge 0$

▸**Page 665** lines 2–4 — 18 Feb 2004

$\left(\frac{y}{N}\right) = 1$ and ... negative Jacobi symbols. $\leadsto$ $\left(\frac{y}{N}\right) \geq 0$ and y is even. Then $y = \bar{x}$, since the other square roots of $\bar{x}^2$ are $N - \bar{x}$ and $(\pm f\bar{x}) \bmod N$; the first of these is odd, and the other two either have negative Jacobi symbols or are simply $\bar{x}$ and $N - \bar{x}$.

▸**Page 666** line −5 of answer 40 — 18 May 2002

$s \leftarrow r \leadsto s' \leftarrow r$

▸**Page 667** line 4 — 21 Jan 1998

$f\left(\frac{N^3}{kp_j}\right) \leadsto f\left(\frac{N^3}{kp_j}, p_{j-1}\right)$

Page 667 last lines of answer 41 — 30 Sep 2009

Lagarias and Odlyzko also developed ... [end of answer] $\leadsto$ Further refinements due to Marc Deléglise, Joël Rivat, Xavier Gourdon, and Tomás Oliveira e Silva have made it possible to deduce that $\pi(10^{23}) = 1925320391606803968923$; see *Revista do DETUA* **4** (2006), 759–768. Lagarias and Odlyzko also developed a completely different approach whereby $\pi(N)$ can be evaluated in $O(N^{1/2+\epsilon})$ steps, using principles of analytic number theory; see *J. Algorithms* **8** (1987), 173–191. But the constant in that O is impracticably large.

▸**Page 668** answer 43 — 14 Feb 2000

line 4: $X_1 \cdots + X_n \leadsto X_1 + \cdots + X_n$
line 23: $2r\epsilon^2 \leadsto 2r\epsilon^{-2}$

Page 669 lines −13 and −12 — 02 Jul 2001

Lecture Notes ... 267–279 $\leadsto$ *J. Cryptology* **13** (2000), 221–244.

▸**Page 670** line 14 of answer 45 — 06 Feb 1998

$x^2 - ay_2 \equiv 1 \leadsto x^2 - ay^2 \equiv 1$

▸**Page 671** line 5 of answer 46 — 09 Feb 1998

(modulo p) $\leadsto$ (modulo $p - 1$)

Page 676 in answer 18 — 30 Jan 2010

[change the step numbers C1, C2, C3, C4 respectively to S1, S2, S3, S4]

▸**Page 676** line 4 of answer 18 — 03 Mar 2000

$u_1v_1 = u_1v_2 \leadsto u_1v_1 = u_2v_2$

▸**Page 677** line 8 — 26 Jan 2001

$UV = 1 \leadsto UV = I$.

▸**Page 677** line 12 — 28 Sep 2002

$T_{(j+1)j} \leadsto T_{(i+1)j}$

Page 677 new answer for exercise 20 — 29 Dec 2008

20. See V. Y. Pan, *Information and Computation* **167** (2001), 71–85.

▸**Page 680** line 3 — 30 Jun 2002

)); see $\leadsto$))); see

▸**Page 681** lines 4–9 of answer 15 — 07 Aug 2001

in (20), or better ... for large p. ↜↝ suggests that $\gcd(x^2-u,(sx+t)^{(p-1)/2}-1)$ will often be a nontrivial factor; and indeed one can show that $(p-1)/2+(0,1,\text{or }2)$ values of s will succeed. In practice, sequential choices seem to work just as well as random choices, so we obtain the following algorithm: "Evaluate $\gcd(x^2-u,x^{(p-1)/2}-1)$, $\gcd(x^2-u,(x+1)^{(p-1)/2}-1)$, $\gcd(x^2-u,(x+2)^{(p-1)/2}-1)$, ..., until finding the first case where the gcd has the form $x+v$. Then $\sqrt{u}=\pm v$." The expected running time (with random s) will be $O(\log p)^3$ for large p.

Page 681 line −8 — 07 Jul 1999

Pub. International Press, to appear) ↜↝ International Press, 1998), 1–11

▸**Page 682** lines 4 and 5 of answer 20 — 18 Feb 2004

$u_j = u_m \sum \alpha_{i_1}\ldots\alpha_{i_{m-j}} \ldots |u_m| \sum \beta_{i_1}\ldots\beta_{i_{m-j}}$ ↜↝ $u_j = \pm u_n \sum \alpha_{i_1}\ldots\alpha_{i_{n-j}}$, an elementary symmetric function, hence $|u_j| \le |u_n| \sum \beta_{i_1}\ldots\beta_{i_{n-j}}$

▸**Page 684** line 2 of answer 21(e) — 03 Feb 2005

$[v]=[w]=2^n$ ↜↝ $[v]^2=[w]^2=2^n$

Page 684 line 2 of answer 22 — 23 Feb 2000

$c\ell(v)$ ↜↝ $c\cdot\ell(v)$

Page 685 line of the last paragraph — 01 Oct 2009

(Göttingen, 1876), 238]. Hensel's name ↜↝ (Göttingen, 1863), 238]. Meanwhile T. Schönemann had independently discovered the lemma and proved uniqueness [*Crelle* **32** (1846), 93–105, §59]. Hensel's name

Page 686 line 2 of answer 26 — 19 May 2005

$D \leftarrow D \vee (D \lll d_j)$, where $\vee$ denotes bitwise "or" and $D \lll d$ denotes D shifted ↜↝ $D \leftarrow D \mid (D \ll d_j)$, where $\mid$ denotes bitwise "or" and $D \ll d$ denotes D shifted

▸**Page 688** line 6 — 06 Jun 1999

$(a+t_2)$ ↜↝ $(\alpha+t_2)$

▸**Page 688** lines 10 and 11 of answer 32 — 13 Jan 1998

However, ζ^p is ... divisor of $f(x)^p$ ↜↝
However, ζ is a root of $g(x^p)$, so $\gcd(f(x),g(x^p))\neq 1$ over the integers, hence $f(x)$ is a divisor of $g(x^p)$. By (5), $f(x)$ is a divisor of $g(x)^p$

▸**Page 691** line 10 of answer 2 — 29 Mar 2000

09 ↜↝ *08*

▸**Page 693** line 4 of answer 15 — 31 Mar 1998

precisely by the ↜↝ precisely by not matching any of the

▸**Page 693** bottom line — 02 Apr 2000

$(j_i\,..\,i_1]$ ↜↝ $(j_1\,..\,i_1]$

▸**Page 695** replacement for lines 2–3 of answer 27 — 25 May 2008

$c(r+1)$ we have $l(n)-\lambda(n)\le r-\lambda(c(r))=l(c(r))-\lambda(c(r))$. The answers for $1\le s\le 7$ are therefore 3, 7, 29, 127, 1903, 65131, 4169527; probably $c(36)$ will require 8.

Page 695 answer 28 — 19 May 2005

line 1: $x \vee y \vee (x + y)$, where "$\vee$" is $\leadsto$ $x \mid y \mid (x + y)$, where "$\mid$" is
line 2: $\nu(x \vee y) + \nu(x \wedge y)$ $\leadsto$ $\nu(x \mid y) + \nu(x \,\&\, y)$

Page 697 replacement for lines 1–5 of answer 37 — 01 Jan 2002

(Solution by D. J. Bernstein.) Let $n = n_m$. First compute 2^e for $1 \le e \le \lambda(n)$, then compute each n_j in $\lambda(n)/\lambda\lambda(n) + O(\lambda(n)\lambda\lambda\lambda(n)/\lambda\lambda(n)^2)$ further steps by the following variant of the 2^k-ary method, where $k = \lfloor \lg\lg n - 2\lg\lg\lg n \rfloor$: For all odd $q < 2^k$, compute $y_q = \sum\{2^{kt+e} \mid d_t = 2^e q\}$ where $n_j = (\ldots d_1 d_0)_{2^k}$, in at most $\lfloor \frac{1}{k} \lg n \rfloor$ steps; then use the method in the final stages of answer 9 to compute $n_j = \sum q y_q$ with at most $2^k - 1$ further additions.

Page 697 line −8 — 31 Jan 2002

proved a comprehensive generalization $\leadsto$ considerably extended

Page 698 new answer — 17 Sep 2005

42. The condition fails at 128 (and in the dual 1, 2, ..., 16384, 16385, 16401, 32768, ... at 32768). Only two reduced digraphs of cost 27 exist; hence $l^0(5784689) = 28$. Furthermore, Clift's programs proved that $l^0(n) = l(n)$ for all smaller values of n.

▸**Page 699** line 2 (the displayed equation) — 28 Jun 1999

$\beta_n)$ $\leadsto$ β_n

▸**Page 699** line 7 of answer 10 — 01 Feb 2005

J. Algebra **5** (1967), 342–357 $\leadsto$ *J. Algebra* **3** (1966), 1–27.

Page 700 lines 9–12 — 10 Aug 2000

J.-C. Lafon ... If all $\leadsto$ Markus Bläser, who showed that $2n^2 + n - 3$ nonscalar multiplications are necessary for $n \ge 2$, and $mn + ns + m - n + s - 3$ in the $m \times n \times s$ case for $n \ge 2$ and $s \ge 2$ [*Computational Complexity* **8** (1999), 203–226]. If all

Page 700 append new sentence to line 4 of answer 14 — 07 Jun 2001

Replace $f_k(s_{n-k+1}, \ldots, s_n, t_1, \ldots, t_{n-k})$ by $f_k(t_1, \ldots, t_{n-k}, s_{n-k+1}, \ldots, s_n)$ in (40) if you prefer to work *in situ*.

▸**Page 701** line 3 of answer 16 — 25 Mar 2001

step C8 $\leadsto$ step T8

▸**Page 703** line 9 of answer 34 — 20 Apr 2000

that uses λ_j $\leadsto$ that uses λ_i

Page 706 line 1 of answer 41 — 11 May 1999

Beware numerical $\leadsto$ Beware of numerical

Page 706 line 6 of answer 41 — 12 Jul 2003

its Applications $\leadsto$ *Its Applications*

▸**Page 707** last line of answer 42 — 28 Jun 1999

180.] $\leadsto$ 180.

Page 708 line −2 of answer 47 — 08 Mar 2000

Ph. D. $\leadsto$ Ph.D.

Page 712 line 2 — 12 Jul 2003

its Applications ↝ *Its Applications*

▸**Page 713** line −4 — 10 Oct 1999

bit-reversal in s'.) ↝ bit-reversal in s'.

▸**Page 714** lines 10 and 11 of answer 60 — 21 Apr 2002

indices of the other. ↝ indices $[k, i, j]$ of the other, and work with an $(mn+sm) \times (ns+mn) \times (sm+ns)$ tensor.

▸**Page 715** line 3 of answer 64 — 28 Jun 1999

$zk3$ ↝ z_{k3}

▸**Page 715** line 3 of answer 66 — 30 Oct 2005

sufficiently large N ↝ sufficiently large n

Page 718 lines 12–15 — 15 Mar 1998

E. Kaltofen ... Of course ... theoretical interest. ↝ Of course ... theoretical interest. E. Kaltofen has shown how to evaluate determinants with only $O\bigl(n^{2+\epsilon}\sqrt{M(n)}\bigr)$ additions, subtractions, and multiplications [*Proc. Int. Symp. Symb. Alg. Comp.* **17** (1992), 342–349]; his method is interesting even with $M(n) = n^3$.

▸**Page 722** line 2 of answer 16 — 22 Apr 1999

$\frac{d}{dt}\bigl(R_{k+1}(t)/V(t)^n\bigr)$ ↝ $\frac{d}{dt}\bigl(R_k(t)/V(t)^{n+1}\bigr)$

Page 722 line 4 of answer 19 — 09 Oct 2010

These coefficients arise ↝ These coefficients, called partial Bell polynomials [see *Annals of Math.* (2) **35** (1934), 258–277], arise

Page 723 line 2 of answer 20 — 22 Aug 2002

Paris **224** (1947) ↝ **224** (Paris, 1947)

Page 724 last line of answer 26 — 11 Jul 1999

to appear ↝ *J. Symbolic Computation* **26** (1998), 339–341

Page 727 Table 2 ..... 06 May 1999

In the next edition I plan to give these constants to 36 hexadecimal places, instead of 45 octal places.

▸**Page 730** entry for Kronecker delta — 14 Aug 1998

1.2.6 ↝ 1.2.3

▸**Page 731** entry for $\Im z$ — 26 Jan 2001

imaginary part z ↝ imaginary part of z

▸**Page 733** entry for $\exp x$ — 09 Aug 1999

1.2.2 ↝ 1.2.9

Page 735 a new appendix — 06 Jan 2011

Beginning with the 26th printing, Volume 2 now has an "Appendix C" analogous to Appendix C of Volume 4A, which lists the page numbers on which named algorithms, theorems, programs, definitions, ..., can be found. A copy of this two-page appendix can be found online in the author's website. The main index therefore begins now on page 737.

Page 736 quotation for bottom of the page — 07 Jan 2011

At any step, arbitrary combinations of algorithms and theorems can be applied to solve a given problem.

— KARSTEN HOMANN and JACQUES CALMET (1995)

Page 735 and following 22 Aug 1997

Miscellaneous changes to the existing index of Volume 2 are collected here, including corrections and amendments to the old entries as well as new entries that are occasioned by the new material. Thus, the lines of the full index that have changed serve also as an index to the present document. However, when a correction or amendment has caused an old index entry to be deleted, the deletion is usually not indicated.

0–1 matrices, 499.
0–1 polynomials, 497, 519, 707.
2-adic numbers, 213, 629.
π (circle ratio), 41, 151, 158, 161, 198, 200, 209, 279–280, 284, 358, 726–727, 733.
  as "random" example, 21, 25, 33, 47, 52, 89, 103, 106, 108, 184, 238, 243, 252, 324–325, 593, 599, 665.
$\rho(n)$ (ruler function), 540.
Addition chains, star, 467, 473–477, 480, 482.
Adleman, Leonard Max, 403, 405, 414, 417, 671.
Agrawal, Manindra (मणीन्द्र अग्रवाल), 396.
Ahrens, Joachim Heinrich Lüdecke, 119, 129–130, 132–134, 136, 137, 140, 141, 588.
al-Bīrūnī, Abū al-Rayḥan Muḥammad ibn Aḥmad (أبو الريحان محمد بن أحمد البيروني), 461.
al-Samaw'al (= as-Samaw'al), ibn Yaḥyā ibn Yahūda al-Maghribī (ابن يحيى بن يهودا المغربي السموءل), 198.
Alexeev, Boris Vasilievich (Алексеев, Борис Васильевич), 117.
Alt, Helmut, 706.
Anderson, Stanley Frederick, 312.
Arithmetic chains, *see* Quolynomial chains.
Armengaud, Joël, 409.
Arney, James W., 385.
Āryabhaṭa I (आर्यभट), 343.
Avanzi, Roberto Maria, 396.
Bell, Eric Temple, polynomials, 722.
Bender, Edward Anton, 385.
Berger, Arno, 262.
Bernoulli, Nicolas (= Nikolaus), 449.
Bernstein, Daniel Julius, 396, 697, 724.
Berrizbeitia Aristeguieta, Pedro José de la Santísima Trinidad, 396.
Bharati Krishna Tirthaji Maharaja, Jagadguru Swami Sri (जगद्गुरु स्वामी श्री भारती कृष्ण तीर्थजी महाराज), 208.
Bhāskara I, Ācārya (भास्कराचार्य), 343.
Bläser, Markus, 700.
Blum, Manuel, integer, 174–176, 183, 416.
Binary recurrences, 318, 466, 634, 692, 714.
Binary search trees, 593.
BINEG computer, 205.
Bolker, Ethan David, 593.
Boolean functions, 173–174.
Boone, Steven Richard, 409.
Booth, Andrew Donald, 608.
Brent, Richard Peirce, 8, 28, 40, 130, 136, 139, 141, 187, 241, 279, 280, 313, 348, 352–353, 355, 356, 382, 386, 403, 501, 529–534, 539–540, 556, 590, 600, 643, 644, 646, 657, 658, 695, 719–721.
Bunimovich, Leonid Abramovich (Бунимович, Леонид Абрамович), 262.
Calmet, Jacques, 736.
Cameron, Michael James, 409.
Canonical signed bit representation, 611.
Cantor, Georg Ferdinand Ludwig Philipp, 209.
Cantor, Moritz Benedikt, 655.
Caramuel de Lobkowitz, Juan, 199–200.
Carmichael, Robert Daniel, numbers, 659, 662.
Cellular automaton, *see* Linear iterative array.
Cervantes Saavedra, Miguel de, 148.
Chapple, Milton Arthur, 530.
Chebotarev, Nikolai Grigorievich (Чеботарёв, Николай Григорьевич), 690.
Cheng, Qi (程岐), 396.
Chinese mathematics, 197–198, 287, 340–341, 486.
Chinese remainder theorem, 285–290, 389, 404, 584.
  for polynomials, 440, 456, 509–510.
Clarkson, Roland Hunter, 409.
Clift, Neill Michael, 477–479, 485.
Cocks, Clifford Christopher, 407.
Codes for difficulty of exercises, xi.
Cohen, Henri José, 345, 658, 687, 712.
Collenne, Joseph Désiré, 201.
Concave function, 125, 139, 245, 627.
Convex function, 125, 139, 245, 684.
Convolution polynomials, *see* Poweroids.
Cooper, Curtis Niles, 409.
Cryptography, 2, 193, 403–407, 415, 417, 505.
David, Florence Nightingale, 3, 566.

Datta, Bibhutibhusan (বিভূতিভূষণ দত্ত) = Bidyāranya, Swami (স্বামী বিদ্যারণ্য), 343, 461.
de La Vallée Poussin, Charles Jean Gustave Nicolas, 381.
Dellac, Hippolyte, 465.
Dewey, Melville (= Melvil) Louis Kossuth, notation for trees, 555.
Dieter, Ulrich Otto, 89, 91, 92, 101, 114, 116, 119, 129–130, 132, 134, 137, 573, 574, 588.
Diophantus of Alexandria (Διόφαντος ὁ Ἀλεξανδρεύς), *see* Diophantine equations.
Dirichlet, Johann Peter Gustav Lejeune, 342.
Discrete Fourier transforms, 169, 305–311, 316–318, 501–503, 506, 512, 516, 520–521, 524, 595.
Dual of an addition chain, 481, 484, 485.
Duffield, Nicholas Geoffrey, 593.
e, as "random" example, 21, 33, 47, 52, 108.
Eckhardt, Roger Charles, 189.
Eichenauer-Herrmann, Jürgen, 32, 558, 559.
Elvenich, Hans-Michael, 409.
ERNIE machine, 3–4.
Euclid's algorithm, 86, 99, 102, 117, 184, 288, 290, 304, 334–337, 340, 579.
Eudoxus of Cnidus, son of Æschinus (Εὔδοξος Αἰσχίνου ὁ Κνίδιος), 335, 359.
Euler, Leonhard (Ейлеръ, Леонардъ = Эйлер, Леонард), xi, 357, 375, 377, 392, 407, 526, 649–651, 653, 655.
Fallacious reasoning, 26, 74, 88, 95–96, 222.
Fast Fourier transform, 73, 306–310, 318, 502, 505, 512, 516, 521, 706, 710, 713–714.
Feijen, Wilhelmus (= Wim) Hendricus Johannes, 636.
Fermat, theorems, 391, 413, 440, 579, 680.
Fibonacci, Leonardo, of Pisa (= Leonardo filio Bonacii Pisano), 197, 208, 280.
  sequence, 27, 37, 213, 264, 360, 468, 483, 660, 666.
Findley, Josh Ryan, 409.
Fine, Nathan Jacob, 90.
Flajolet, Philippe Patrick Michel, 355, 366, 449, 541, 644.
Fluhrer, Scott, 542.
FORTRAN language, 188, 193, 279, 600, 602.
Fowler, Thomas, 208.
Frobenius, Ferdinand Georg, 539, 681, 689.
Gathen, Joachim Paul Rudolf von zur, 449, 611, 673, 687.
Gauß (= Gauss), Johann Friderich Carl (= Carl Friedrich), 20, 101, 363, 417, 422, 449, 578, 679, 685, 688, 701.
Gerke, Friedrich Clemens, 653.
Gimeno Fortea, Pedro, 187.
Goldstein, Daniel John, 593.
Goodman, Allan Sheldon, 108.
Goulard, Achille, 477.
Gourdon, Xavier Richard, 355, 449, 667.
Granville, Andrew James, 396, 659.
Gray, Frank, binary code, 209, 568, 699.
Gray levels, multiplication of, 284.
Greek mathematics, 196–197, 225, 335–337, 359.
Hadamard, Jacques Solomon, transform, 173, 502.
Hajratwala, Nayan (ઘયન ખજરતવાલા), 409.
Hensley, Douglas Austin, 366, 373.
Hilferty, Margaret Mary, 134.
Hill, Theodore Preston, 262.
Hindu mathematics, 197, 208, 209, 281, 287, 343, 387, 461, 648.
Hofstadter, Douglas Richard, 330.
Homann, Karsten, 736.
Hurwitz, Adolf, 345, 375, 376, 649.
Ibn Ezra (= Ben Ezra), Abraham ben Meir (אברהם בן מאיר אבן עזרא), also known as Abū Isḥaq Ibrāhim al-Mājid (أبو إسحق إبراهيم الماجد بن عزرا), 197.
IEEE standard floating point, 226, 246, 602.
in situ transformation, 700.
Indian mathematics, 197, 208, 209, 281, 287, 343, 387, 461, 648.
Inverse modulo 2^e, 213, 629.
Inverse modulo m, 26, 354, 445, 456, 646.
Jacobi symbol, 413–414, 415, 655, 662, 668.
Jensen, Geraldine Afton, 466.
Kayal, Neeraj (नीरज कयाल), 396.
Killingbeck, Lynn Carl, 103, 107.
Kornerup, Peter, 332–333, 629, 657.
Krandick, Werner, 625, 629.
Kurowski, Scott James, 409.
Lagrange (= de la Grange), Joseph Louis, Comte, 375, 378, 456, 503, 527, 533, 549, 649, 653, 655.
Lakshman, Yagati Narayana (ಲಕ್ಷ್ಮಣ ಯಾಗಟಿ ನಾರಾಯಣ), 455.
Lalescu, Liviu, 186.
Lebesgue (= Le Besgue), Victor Amédée, 341, 662.
Léger, Roger, 587.
Lehmer, Derrick Norman, 278, 661.
Lenstra, Arjen Klaas, 118, 403, 417, 453, 712.
Leva, Joseph Leon, 132.
Levine, Kenneth Allan, 104.
Lochs, Gustav, 372–373.
Loewenthal, Dan (דן לבנטל), 291.
Lund, Carsten, 593.
Machine language versus higher-level languages, 16, 185.
Maiorana, James Anthony, 557.
Marczyński, Romuald Władysław, 205.

Marsaglia, George, 3, 23, 29, 33, 40, 47, 62, 71, 72, 75, 78, 108, 114–115, 119, 122, 123, 128, 133–135, 179, 544, 546–547, 549, 551, 565, 588.
Masking, 322, 328–329, 389–390, 671.
Matthew, Saint (Ἅγιος Ματθαῖος ὁ Εὐαγγελιστής), 735.
Maze, Gérard, 645.
Metropolis, Nicholas Constantine (Μητρόπολης, Νικόλαος Κωνσταντίνου), 4, 189, 240, 242, 327.
Mignotte, Maurice, 450, 683.
Mihăilescu, Preda-Mihai, 396.
Miller, James (= Jimmy) Milton, 108.
Minus zero, 202, 244–245, 249, 268, 274.
mod m arithmetic, division, 354, 445, 499; *see also* Inverse modulo m.
Moews, David John, 593.
Monte Carlo, 2, 29, 55, 114, 185, 189.
Morley, Frank Vigor, 199.
Moses, Joel, 454–455.
Motzkin, Theodor (= Theodore) Samuel (תיאודור שמואל מוצקין), 378, 490, 494, 495, 497, 518, 519, 705.
Multiple-precision constants, 352, 362, 366, 384, 659, 663, 712, 726–728.
Multiplication, two's complement, 608.
Multiprimality: Total number of prime factors, 384.
Nandi, Salil Kumar (सलिल कुमार नन्दी), 278.
Nārāyaṇa Paṇḍita, son of Nṛsiṁha (नारायण पण्डित, नृसिंहस्य पुत्र:), 387.
Needham, Noel Joseph Terence Montgomery (李約瑟), 287.
Neumann, John von (= Margittai Neumann János), 1, 3–4, 26, 36, 119, 125, 128, 138, 140, 202, 226, 278, 327.
Newton, method for rootfinding, 278–279, 312, 486, 529, 629, 719.
Nicomachus of Gerasa (Νικόμαχος ὁ ἐκ Γεράσων), 659.
Nowak, Martin R., 409.
Nussbaumer, Henri Jean, 521, 710.
Nystrom, John William, 201.
Oliveira e Silva, Tomás António Mendes, 386, 667.
Organ-pipe order, 378.
Packing, 109.
Palmer, John Franklin, 222.
Pan, Victor Yakovlevich (Пан, Виктор Яковлевич), 490, 492, 497, 500, 505, 507, 515, 517, 519, 521, 677, 699, 703, 705, 706, 714, 715, 721.
Pemantle, Robin Alexander, 542.
Percival, Colin Andrew, 632.
Pi (π), 41, 151, 158, 161, 198, 200, 209, 279–280, 284, 358, 726–727, 733.
  as "random" example, 21, 25, 33, 47, 52, 89, 103, 106, 108, 184, 238, 243, 252, 324–325, 593, 599, 665.
Pigeonhole principle, 286.
Piṅgala, Ācārya (आचार्य पिङ्गल), 461.
Planck, Karl Ernst Ludwig Marx (= Max), constant, 214, 227, 238, 240.
Polynomial, resultant, 433, 674, 690.
Potency, 24–26, 36, 47, 52, 73, 83, 87–88, 92, 105, 184.
Powers, Donald (= Don) Michael, 312.
Powers, Ralph Ernest, 396, 407.
Priority sampling, 148.
Quolynomial chains, 498, 524, 704–705.
Rackoff, Charles Weill, 179.
Raleigh (= Ralegh), Walter, 199.
Random integers in a bounded set, 2–3, 6–7, 119–121, 138, 162–163, 185–186.
Random walk test, 34, 79.
Reciprocals, mod 2^e, 213, 629.
Rehkopf, Donald Caspar, 41.
Reitwiesner, George Walter, 213, 280.
Rejection method, 125–126, 128–129, 134, 138, 139, 591.
Reservoir sampling, 143–144, 147, 148.
Resultant of polynomials, 433, 674, 690.
Robbins, David Peter, 593.
Rosińska, Izabela Grażyna, 198.
Rozier, Charles Preston, 324.
RSA box, 404, 406.
Ruler function $\rho(n)$, 540.
Saxena, Nitin (नितिन सक्सेना), 396.
Scholz–Brauer conjecture, 478–479, 485.
Schönemann, Theodor, 457, 685.
Schönhage, Arnold, 292, 302–303, 305, 306, 311, 317, 328, 470, 484, 500, 522, 629, 638, 656, 672, 696, 715.
Schönhage–Strassen algorithm, 306–311, 317.
Schubert, Friedrich Theodor von, 450.
Selfridge, John Lewis, 394, 396, 412, 665.
Seneschal, David, 589.
Seroussi Blusztein, Gadiel (גדיאל סרוסי), 712.
Shafer, Michael William, 409.
Shamir, Adi (עדי שמיר), 403, 405, 416, 505, 599, 669.
Shokrollahi, Mohammad Amin (محمد امین شکرالهی), 515.
Shukla, Kripa Shankar (कृपा शंकर शुक्ला), 208, 648.
Sideways addition, 463, 466.
Sierpiński, Wacław Franciszek, 666.
Signed magnitude representation, 202–203, 209–210, 247, 266.
Sikdar, Kripasindhu (कृपासिन्धु शिकदार), 327.
Silverman, Joseph Hillel, 402.
Singh, Parmanand (परमानन्द सिंह), 387.
SLB (shift left rAX binary), 339, 340.
Smirnov, Nikolai Vasilievich (Смирнов, Николай Васильевич), 57.

Smith, Edson McIntyre, 409.
Smith, Robert LeRoy, 228.
Squares, sum of two, 579–580.
SRB (shift right rAX binary), 339, 340, 481.
Stahnke, Wayne Lee, 31.
Star chains, 467, 473–477, 480, 482.
Steele, Guy Lewis, Jr., 635–636, 638.
Ştefănescu, Doru, 450.
Stern, Moritz Abraham, 654.
Strindmo, Odd Magnar, 409.
Sturm, Jacques Charles François, 434, 438, 674.
Subbarao, Mathukumalli Venkata (మతుకుమల్లి వెంకట సుబ్బారావు), 469.
Subnormal floating point numbers, 246.
Sugunamma, Mantri (மந்திரி சுகுணம்மா), 469.
Szekeres, György (= George), 570.
Tarski (Tajtelbaum), Alfred, 718.
Tate, John Torrence, Jr., 402.
Thorup, Mikkel, 593.
Toeplitz, Otto, system, 721.
Totient function $\varphi(n)$, 19–20, 289, 369, 376, 583, 646.
Trabb Pardo, Luis Isidoro, 661.
Trinomials, 32, 40, 572.
Turán, Pál (= Paul), 372, 649.
Two's complement notation, 15, 188, 203–204, 228, 275–276, 608.
Tydeman, Frederick John, 638.
Ulam, Stanislaw Marcin, 138, 140, 189.
Vallée, Brigitte, 352, 355, 366, 644, 645.
Vitányi, Pál Mihály (= Paul Michael) Béla, 179.
von Neumann, John (= Margittai Neumann János), 1, 3–4, 26, 36, 119, 125, 128, 138, 140, 202, 226, 278, 327.
von Schubert, Friedrich Theodor, 450.
von zur Gathen, Joachim Paul Rudolf, 449, 611, 673, 687.
Vuillemin, Jean Etienne, 629, 649.
Walker, Alastair John, 120, 127, 139.
Wattel, Evert, 466.
Welford, Barry Payne, 232.
White, Jon L, 635–636, 638.
Williams, Hugh Cowie, 380, 390, 394, 401, 412, 415, 661, 664.
Winograd, Shmuel, 280, 316, 500, 501, 507, 509, 512–514, 520, 523, 705–707, 710, 712, 714.
WM1 (word size minus one), 252, 267, 613.
Wolff von Gudenberg, Jürgen Freiherr, 242.
Wunderlich, Charles Marvin, 390, 394, 399–400.
XOR (bitwise exclusive-or), 31, 32, 193, 419.
Zero, minus, 202, 244–245, 249, 268, 274.