

The Cause of Brownian Motion

Liu, Jerry Z. Ph.D.

ZJL@CS.Stanford.EDU

Stanford University, California, USA

Abstract

For the past 200 years, the Brownian motion of small particles has been observed and studied due to its significance in understanding physics at the microscopic scale. Albert Einstein statistically modeled the behavior of these particles, but the mechanism driving Brownian motion has remained unclear. It was previously suggested that molecules push the particles in Brownian motion. However, this study demonstrates that the random movement of molecules cannot provide sufficient momentum to drive Brownian motion. Instead, we propose that Brownian motion is driven by transimpact—an interaction between atoms and a side effect of atomic electron transitions. When energy is absorbed, an electron is excited to a higher orbital, increasing the volume of the atom and reducing the distance to adjacent particles. This disrupts the existing spatial arrangement between atoms and alters the balance of attractive and repulsive forces. The transition induces a repulsive force between particles, with an explosive impact that pushes the particles apart. Transimpact may indeed be the fundamental interaction behind many physical processes and phenomena, such as particle motion, phase transitions, and the rotation of Crookes radiometers. Additionally, transimpact facilitates the exchange of internal energy between a system's potential and kinetic energy, as well as among different components, thereby maintaining equilibrium.

Introduction

In 1827, while studying the pollen grains of the plant *Clarkia pulchella* suspended in water under a microscope, Robert Brown observed the jittery motion of minute particles.^[1-5] By repeating the experiment with particles of inorganic matter, he ruled out the possibility that the motion was life-related. This motion was later named after him as Brownian motion. Since then, it has been recognized that Brownian motion is not unique to pollen but is a common phenomenon observable in daily life. Most people might have noticed dust particles dancing in a beam of light in a dark room. Indeed, Brownian motion reflects a fundamental interaction between particles at a microscopic scale in physics.

In 1905, Albert Einstein statistically modeled the movement of particles in Brownian motion^[6-8] providing a method to indirectly demonstrate the existence of atoms and molecules, which were not well established at the time. While it was postulated that the random movement of fluid molecules pushes the particles in Brownian motion, the underlying physics governing this motion remains unclear to this day. Can the randomly moving molecules have sufficient momentum to drive the particles in Brownian motion? If not, what force pushes the particles? These questions may relate to some of the fundamental processes underlying thermodynamics and many other physical phenomena, which is the primary motivation of this study.

Insufficient Momentum from Molecule Movement

To estimate the velocity required for molecules to drive Brownian motion, let us consider a grain of pollen initially at rest. It is struck by a water molecule moving at velocity v . To maximize the estimated momentum transferred from the molecule, let's assume the water molecule is bounced back at the same velocity in the opposite direction, and the momentum is completely transferred to the pollen. The mass of a water molecule is approximately 2.99×10^{-26} kg. The change in momentum of the water molecule before and after the collision is $v \times 2 \times 2.99 \times 10^{-26}$ or $v \times 5.98 \times 10^{-26}$.

The *Clarkia pulchella* pollen is about 5 microns long (50×10^{-6} m). To be conservative regarding the pollen's size, let's assume a pollen grain to be cubic with a side length of 1×10^{-6} m, corresponding to a volume of 1×10^{-18} m³. Since the pollen grains are suspended in the water, their density should be approximately the same as that of water (997 kg/m³ at 25 °C). Assuming that the pollen picks up a velocity of 1 cm/s (0.01 m/s), the lower range of pollen's motion, after the collision, it will gain a momentum of 9.97×10^{-18} kg•m/s. This momentum should equal the momentum change of the water molecule:

$$(1) \quad v \times 5.98 \times 10^{-26} = 9.97 \times 10^{-18}$$

So, to push the pollen at a velocity of 1 cm/s, the velocity of the water molecule must be greater than

$$(2) \quad v = 1.67 \times 10^8 \text{ m/s}$$

This velocity exceeds 56% of the speed of light ($\sim 3 \times 10^8$ m/s), which is already an unrealistic velocity for water molecules. For a conservative estimate in our initial calculation, we used a small cubic pollen grain with a volume of 1×10^{-18} . However, for a more realistic estimation, consider a typical *Clarkia pulchella* pollen grain with dimensions 5×10^{-5} m in length, 4×10^{-5} m in width, and 4×10^{-5} m in thickness, or a volume of 8×10^{-14} m³. To move such pollen, a water molecule would need to travel at a velocity greater than 1.34×10^{13} m/s, which is faster than the speed of light. The average velocity of water molecules, based on the kinetic energy distribution at a temperature of 20 °C, is approximately 590 m/s.^[9]

This estimate aligns with the speed of sound in water (1500 m/s),^[10] which serves as the upper velocity bound for water molecules since sound waves are propagated by these molecules. The velocity of water molecules can also be compared to that of air molecules. The average velocity of air molecules is less than 1,200 m/s. Given that water has a much higher viscosity than air, the average velocity of water molecules should be substantially slower than that of air molecules. The estimated velocity of water molecules required for Brownian motion is multiple orders of magnitude higher than what is realistically possible. Thus, the momentum of a single water molecule is insufficient to push pollen in Brownian motion.

One might argue that the synchronous impacts of multiple water molecules could collectively push the pollen. To move a typical pollen grain with a volume of 8×10^{-14} m³ at a velocity of 1 cm/s, with the average velocity of water molecules (590 m/s), we need 2.26×10^{10} molecules to strike a pollen surface simultaneously. This number, N , is estimated based on the following equation:

$$(3) \quad N \times 590 \times 5.98 \times 10^{-26} = 0.01 \times 997 \times 8 \times 10^{-14}$$

To estimate this possibility, consider the large surface area of the typical pollen grain, approximately $5 \times 4 \times 10^{-10} \text{ m}^2$, being struck by all the water molecules on this side at the same time. Based on the density of water (997 kg/m^3) and the mass of a water molecule ($2.99 \times 10^{-26} \text{ kg}$), the number of water molecules per cubic meter is approximately 3.33×10^{28} , which translates to 1.04×10^{19} molecules per square meter. Therefore, the maximum number of water molecules on this surface should be around 2.08×10^{10} , which is less than the required 2.26×10^{10} molecules needed to move the pollen. Thus, even the collective impact of all the water molecules is insufficient to provide the necessary momentum to move the pollen.

Furthermore, even if the surface area of the pollen could accommodate a sufficient number of water molecules, all the molecules on the opposite side would have to move away from the pollen for it to be displaced. Additionally, due to the random nature of these collisions, molecules typically come from different directions and tend to cancel each other out. The likelihood of multiple molecules acting collectively is extremely low. Consequently, the collective argument for Brownian motion lacks support. If the movement of molecules cannot provide sufficient momentum for Brownian motion, what mechanism drives it?

The Mechanism Driving Brownian Motion

"Transimpact", a term derived from "transition impact", refers to an interaction between atoms resulting from a side effect of atomic electron transitions.^[11] Typically, attractive forces are induced between atoms or molecules that are close. When the space between them becomes too small, the repulsion of their valence electrons increases rapidly. At a certain distance, a balance is achieved between these repulsive and attractive forces. This balance is often disrupted by atomic electron transitions. When an electron absorbs energy, such as from electromagnetic waves, it will be excited to a higher orbital, usually within a few nanoseconds.^[12-13] As a result, the electron cloud and the atom's volume expand, reducing the distance to adjacent atoms. This sudden reduction in space disrupts their balance and significantly increases the repulsion between them. The impact is explosive and can exert a substantial force on adjacent particles.




Figure 1: Process of transimpact.

As illustrated in Figure 1A, two single-atom molecules are positioned close to each other, with their attraction and repulsion balanced at a small separation. Upon absorbing energy, an electron in one of the molecules undergoes an atomic orbital transition, increasing the molecule's size and reducing the space between them, as shown in Figure 1B. This expansion

increases the repulsion between the molecules, pushing them apart and causing them to move in opposite directions, as depicted in Figure 1C.^[11,14]

To illustrate this interaction, let us consider the electron orbital transition of an oxygen atom. An atomic radius is primarily determined by its valence orbital. An oxygen atom has 8 electrons, with a ground state configuration of $1s^2 2s^2 2p^4$, giving it a radius of approximately 48 pm. In its typical excited state, the configuration is $1s^2 2s^2 2p^3 3s^1$, with a radius around 118 pm. Thus, when an oxygen atom becomes excited, its radius more than doubles, leading to a volume increase of over 1,200% within a few nanoseconds. This expansion effect is likely even more pronounced in hydrogen atoms. This rough estimate demonstrates how a water molecule could expand significantly as it absorbs energy from its surroundings.

The impact of transimpact can be visualized using the burst of popcorn. Consider the momentum exerted on adjacent objects by the popcorn as it pops. This analogy is similar to the interaction that occurs when an electron in a water molecule becomes excited next to a pollen grain. It is believed that this mechanism drives the Brownian motion of pollen in water.

Since atomic electron transitions are common interactions at the microscopic scale, transimpact is likely a fundamental interaction that influences many physical processes. The particles or objects affected by transimpacts can include other atoms, molecules, pollen, dust particles, or even the walls of a container. When a transimpact occurs between two molecules, the molecules are pushed apart and gain kinetic energy, manifesting as vibration in solids or random motion in fluids. In other words, transimpacts facilitate the transformation of energy from potential to kinetic, or kinetic energy derived from potential energy. We will explore some thermodynamic processes related to transimpacts in the following sections.

The Force Driving Crookes Radiometers

Transimpact provides a convincing explanation for the operation of Crookes radiometers.^[14] The vanes of a Crookes radiometer rotate when exposed to light. Despite its long history, the mechanism behind its rotation remained a mystery for over a century. Many theories were proposed, but none could account for all observations. When a transimpact occurs between an air molecule and the vane of the radiometer, the vane is pushed by the impact. Under light exposure, the black side of the vanes absorbs energy more effectively, resulting in a higher frequency of transimpacts compared to the white side. This imbalance in impacts causes the vanes to rotate. Observations from extensive light experiments have provided compelling evidence supporting this theory.

Most existing theories attribute the rotation of the vanes to airflow or differences in air pressure inside the radiometer. For these theories to be valid, the air within the radiometer would need to be heated to create pressure differentials, a process that takes time. For the vanes to start rotating, the driving force must be strong enough to overcome the friction between the vanes and the mounting spindle. Consequently, the vanes cannot move initially until this friction is overcome. Furthermore, even after rotation begins, acceleration should start from zero and gradually increase, as it is proportional to the net driving force and should rise as the air heats up. However, this initial phase of zero or low acceleration is absent in nearly all observations.

In contrast, the transimpact theory predicts that acceleration begins at its maximum because atomic electron transitions occur as soon as the radiometers are exposed to light. As the rotation gains speed, air resistance increases, which

reduces the driving force and, consequently, the acceleration. Therefore, the rotation speed should initially increase until it reaches a peak, where the driving force is balanced by air resistance. This pattern is observed in almost all experiments. To further verify this prediction, an experiment was conducted using a powerful flashlight. The speed changes observed during the experiment are compiled in Table 1. Each acceleration value in the table represents the average between two sample points of the rotation speed.

Time (s)	0	2	5	17	30	43	64	78	100	110	120	128
Speed (Hz, rev/s)	0	7.5	15	22.5	25	22.5	15	7.5	3.75	3	3	0
Acceleration (Hz/s)		3.75	2.5	0.625	0.192	-0.192	-0.357	-0.536	-0.17	-0.075	0	-0.375

Table 1: Changes in radiometer speed and acceleration during the flashlight experiments.




Figure 2: Changes in radiometer speed and acceleration during the flashlight experiment.

The experimental data presented in Table 1 are plotted in Figure 2. The striking agreement between the observations and predictions provides compelling evidence for the validity of the transimpact theory in explaining the behavior of Crookes radiometers.

Misconception about Temperature Representation

It is commonly believed that temperature represents the average kinetic energy of particle motion in substances. This misconception will be clarified by examining the role of transimpacts in particle interactions. Transimpacts cause particle vibrations in solids and motion in fluids. In gases, this motion becomes so pronounced that it leads to the belief that the kinetic energy of particle motion is the primary component of thermal energy, which temperature represents. This misconception is further reinforced by the empirical law of ideal gases. Based on this law, the kinetic theory of gases was developed, establishing a correlation between temperature and the average speed of gas molecules:

$$(4) \quad v = \sqrt{\frac{3k_b T}{m}},$$

where m denotes the mass of a gas molecule, T indicates the absolute temperature, k_b represents the Boltzmann constant ($1.38 \times 10^{-23} \text{ J/K}$), and v is the predicted average speed of the gas molecules.^[15-21] Therefore, the average kinetic energy due to the motion of molecules in an ideal gas can be derived from the kinetic theory:

$$(5) \quad K = \frac{1}{2}mv^2 = \frac{3}{2}k_b T,$$

which suggests the kinetic energy K in the system is proportional to the absolute temperature of the system. This equation reflects the common belief that temperature represents the average kinetic energy in a system. If kinetic energy were the sole determinant of a system's temperature, changes in temperature and kinetic energy could be used to calculate specific heat. Consequently, the specific heat of ideal gases can be predicted using the kinetic theory of gases:

$$(6) \quad C_k = \frac{K}{mT} = \frac{3k_b}{2m}$$

which corresponds to a prediction of (transitional) isometric molar heat capacity:

$$(7) \quad C_v = \frac{3R}{2}$$

where R denotes the ideal gas constant ($8.31 \text{ J/mol}\cdot\text{K}$). This prediction closely matches measurements for gases with monatomic molecules with an accuracy of 99% or higher. However, the kinetic theory of gases fails to accurately predict the specific heat for gases with polyatomic molecules. The predicted values for gases with polyatomic molecules are consistently lower than the measured values. The prediction error appears to be roughly proportional to the number of bonds in the gas molecules, as illustrated in Figure 3.^[22]




Figure 3: The correspondence between unaccounted molar-specific heat in the kinetic theory and the number of bonds. The unaccounted molar-specific heat is measured by offsetting the molar-specific heat predicted by the kinetic theory. The slope of the fitted trendline is about 6.56 with an R^2 of 0.98.

For instance, the predicted specific heat for octane is only 7% of the measured value, indicating a 93% prediction error. In other words, only 7% of the specific heat energy is allocated to increasing the kinetic energy for molecular motion. Numerous adjustments have been introduced to the theory, such as the equipartition of kinetic energy among vibrational, rotational, and translational motions, and considerations of the degrees of freedom in complex molecules.^[23-30] However, these adjustments have all failed to account for the specific heat discrepancies shown in Figure 3.

The root cause of this discrepancy lies in the misconception that kinetic energy solely represents temperature. In reality, temperature measures thermal energy, which encompasses both kinetic and potential energy. Potential energy can be stored between particles under the influence of certain forces, such as chemical bonds. Bonds act like springs for energy storage: when work, such as specific heat, is done to expand a bond's space, the additional energy is stored in the extended bond. Similarly, energy can be stored in an excited electron in higher orbitals. When the electron returns to a lower orbital or the bond contracts, the stored energy is released as electromagnetic waves, which may be absorbed by other electrons or bonds. Thermal energy is this dynamic portion of internal energy and is typically measured by thermometers as temperature. This explains the correlation between the mispredicted specific heat and the number of bonds in gas molecules, as shown in Figure 3.

A transimpact occurs in every potential energy exchange process between different components, where a fraction of the potential energy is converted into kinetic energy for particle motion. Thus, kinetic energy is derived from potential energy and mirrors its changes, creating the delusion of a direct correlation with temperature. However, this relationship is indirect, mediated by potential energy, and will break down during a phase transition at a constant temperature, as discussed next. Consequently, temperature is primarily influenced by potential energy. This becomes clearer when considering the nature of absolute zero temperature, as kinetic energy is not zero even at absolute zero.^[31]

The Cause of Phase Transition

Transimpact may also be the true cause of phase transitions.^[32] A common textbook misconception attributes phase transitions to the increasing intensity of molecular vibrations at high temperatures. Phase transitions are influenced by the interplay of three forces within a system: pressure and bond forces that hold molecules together, and transimpacts that tend to break them apart. The balance among these three forces determines the state of matter. For instance, molecules in a solid can be held together because

$$(8) \quad T < B + P$$

where T denotes the average transimpact force, B indicates the bond force, and P stands for the pressure. A phase transition occurs when

$$(9) \quad T > B + P$$

The impact of transimpacts is more intense than that of molecular vibrations, as kinetic energy is derived from transimpacts. Bonds are likely to be broken by transimpacts before vibrations reach such high intensity. This theory also provides a better explanation for phase transitions occurring at constant temperatures.

This relates to another common textbook misconception, which posits that since the energy in a phase transition is used to break bonds between molecules, [there's no net increase in the particles' kinetic energy](#), and therefore, no rise in temperature. An implicit assumption in this notion is that temperature represents the average kinetic energy. However, as discussed earlier, this is not accurate. Temperature primarily measures the emission level related to the potential energy of a system.

In the case of a phase transition, such as from ice to water, molecules are liberated from their fixed positions within hydrogen bonds, and their kinetic energy must increase through the process. Consequently, if temperature accurately reflected the average kinetic energy of a system, it would be expected to rise during the transition. However, the temperature remains constant during the transition. This contradiction arises because temperature is not directly associated with kinetic energy but with potential energy.

During a phase transition, the temperature remains constant because it is constrained by the potential energy limits of hydrogen bonds until the bonds break. As latent heat is added to the system, the bonds expand to accommodate more potential energy. Simultaneously, the strength of the bonds decreases as they are stretched, and the average intensity of transimpacts increases, making the bonds more vulnerable. Eventually, the phase transition occurs as the bonds are broken by transimpacts. The potential energy of these bonds is limited by the intensity of transimpacts that the bonds can withstand.

As demonstrated earlier, the energy exchange between different components occurs through transimpacts, maintaining equilibrium, particularly in their emission levels. If one component has a higher level of potential energy, the exchange will eventually balance it to a common level, and vice versa. Essentially, the emission level of the entire system is anchored by the potential energy limits of the bonds. Therefore, because the system's potential energy level is fixed, the temperature, measured by the emission level of the bonds, remains constant during a phase transition.

Conclusions

Transimpact is the underlying mechanism behind many physical interactions and phenomena, including the vibration of molecules in solids, the random motion of molecules in fluids, the Brownian motion of small particles, the rotation of Crookes radiometers, and phase transitions, among others. The random movement of molecules does not have sufficient momentum to drive Brownian motion; instead, Brownian motion is driven by transimpact. Temperature is typically a measure of the average potential energy level, rather than the kinetic energy of a system. Different forms of energy can be exchanged through transimpacts to maintain a certain balance within a system. These equilibrium processes create the delusion of a direct correlation between temperature and kinetic energy.

See Also

- [The Second Law of Thermodynamics is not Universal \(PDF\)](#)
- [The Cause of Brownian Motion \(PDF\)](#)
- [The Process Driving Crookes Radiometers \(PDF\)](#)
- [The Nature of Absolute Zero Temperature \(PDF\)](#)

- [Can Temperature Represent Average Kinetic Energy? \(PDF\)](#)
- [Why Phase Transition Temperature Remains Constant \(PDF\)](#)
- [Is Thermal Expansion Due to Particle Vibrating? \(PDF\)](#)
- [Misconceptions in Thermodynamics \(PDF\)](#)
- [Is There a Sea of Free Electrons in Metals? \(PDF\)](#)
- [Unified Theory of Low and High-Temperature Superconductivity \(PDF\)](#)
- [Superfluids Are Not Fluids \(PDF\)](#)
- [Electron Tunnel \(PDF\)](#)
- [Superconductor Origin of Earth's Magnetic Field \(PDF\)](#)
- [Tidal Energy Is Not Renewable \(PDF\)](#)
- [The Simplest Derivation of \$E = mc^2\$ \(PDF\)](#)
- [How to Understand Relativity \(PDF\)](#)
- [The Evolution from the Law of Gravitation to General Relativity \(PDF\)](#)
- [Science vs. Mathematics \(PDF\)](#)
- [Potential Problems of AI-Created Content \(PDF\)](#)
- [DeepSeek vs. ChatGPT at Advanced Levels \(PDF\)](#)

Revision History

- [06/16/2019: Initial Post of This Article.](#)
- [06/03/2021: Revision for Grammar Check.](#)
- [01/07/2023: Adding More Evidence and Removing Energy Theory Sections.](#)
- [10/01/2023: Revision for Clarity and Section Titles.](#)
- [05/27/2024: Adding Examples of Processes Related to Transimpacts.](#)
- 06/30/2024: Updating for Synchronized Motion of Multiple Molecules.

References

1. Feynman, R. (1964). "[The Brownian Movement](#)". The Feynman Lectures of Physics, Volume I. pp. 41-, 1.
2. Li, T.; et al. (2010). "[Measurement of the instantaneous velocity of a Brownian particle](#)". *Science*. **328** (5986): 1673–1675. doi:[10.1126/science.1189403](#). PMID [20488989](#).
3. Weiss, G.H. (1994). "*Aspects and applications of the random walk*". North Holland.
4. Ben-Avraham, D.; Havlin, S. (2000). "*Diffusion and reaction in disordered systems*". Cambridge University Press.
5. Morozov, A. N.; Skripkin, A. V. (2011). "Spherical particle Brownian motion in viscous medium as non-Markovian random process". *Physics Letters A*. **375** (46): 4113–4115. doi:[10.1016/j.physleta.2011.10.001](#).
6. Einstein, A. (1905). "[Über die von der molekularkinetischen Theorie der Wärme geforderte Bewegung von in ruhenden Flüssigkeiten suspendierten Teilchen](#)" [On the Movement of Small Particles Suspended in Stationary Liquids Required by the Molecular-Kinetic Theory of Heat] (PDF). *Annalen der Physik* (in German). **322** (8): 549–560. [Bibcode:1905AnP...322..549E](#). doi:[10.1002/andp.19053220806](#). [Archived](#) (PDF) from the original on 9 October 2022.
7. Einstein, A. (1926). "[Investigations on the Theory of the Brownian Movement](#)". Dover Publications.

8. Stachel, J. (1989). "[*Einstein's Dissertation on the Determination of Molecular Dimensions*](#)". The Collected Papers of Albert Einstein, Volume 2. Princeton University Press.
9. Vertical Learning Labs. (2011). "[Molecular Speed Distribution](#)". Introduction to Chemistry, Unit 1.5.
10. Nieu Kirk S. (2022). "[Understanding Ocean Acoustics](#)". NOAA.
11. Liu, J.Z. (2022), "[Misconceptions in Thermodynamics](#)", Stanford University. Archived ([PDF](#)).
12. Vijay, R.; et al. (2011). "Observation of Quantum Jumps in a Superconducting Artificial Atom". *Physical Review Letters*. **106** (11): 110502. arXiv:[1009.2969](#). doi:[10.1103/PhysRevLett.106.110502](#). PMID [21469850](#).
13. Itano, W.M.; et al. (2015). "[Early observations of macroscopic quantum jumps in single atoms](#)". *International Journal of Mass Spectrometry*. **377**: 403. doi:[10.1016/j.ijms.2014.07.005](#).
14. Liu, J.Z. (2019), "[The Process Driving Crookes Radiometers](#)", Stanford University. Archived ([PDF](#)).
15. [Krönig, A.](#) (1856). "[Grundzüge einer Theorie der Gase](#)". *Annalen der Physik und Chemie* (in German). **99** (10): 315–22. [Bibcode:1856AnP...175..315K](#). doi:[10.1002/andp.18561751008](#). [Facsimile at the Bibliothèque nationale de France \(pp. 315–22\)](#).
16. [Clausius, R.](#) (1857). "[Ueber die Art der Bewegung, welche wir Wärme nennen](#)". *Annalen der Physik und Chemie* (in German). **176** (3): 353–79. [Bibcode:1857AnP...176..353C](#). doi:[10.1002/andp.18571760302](#). [Facsimile at the Bibliothèque nationale de France \(pp. 353–79\)](#).
17. Tuckerman, M.E. (2010). "Statistical Mechanics: Theory and Molecular Simulation". (1st ed.). P. 87. [ISBS 978-0-19-852526-4](#).
18. De Podesta, M.; et al. (2013). "A low-uncertainty measurement of the Boltzmann constant". *Metrologia*, **50** (4): S213-S216, BIPM & IOP Publishing Ltd.
19. Maxwell, J.C. (1875). "The Question of the Atom". *J. Chem. Soc. (London)*, Facsimile published in Mary Jo Nye, **1875** (28): 493-508.
20. Reif, F. (1965). "Fundamentals of Statistical and Thermal Physics". *McGraw-Hill*, New York.
21. Carey, V. (1999). "Statistical Thermodynamics and Microscale Thermophysics". Cambridge University Press.
22. Liu, J.Z. (2023), "[Can Temperature Represent Average Kinetic Energy?](#)", Stanford University. Archived ([PDF](#)).
23. Boltzmann, L. (1895). "On certain Questions of the Theory of Gases". *Nature*, **51** (1322): 413-415.
24. Planck M. "On the Law of the Energy Distribution in the Normal Spectrum". *Ann. Phys.*, **4** (553): 1-11.
25. Einstein, A. and Stern, O. (1913). "Some Arguments for the Assumption of Molecular Agitation at Absolute Zero". *Ann. Phys.*, **40** (551) 551-560.
26. Thomson, W. (1904). "Historical and Philosophical Perspectives". *Baltimore Lectures*. Baltimore: Johns Hopkins University Press. Sec. 27.
27. Rayleigh, J.W.S. (1900). "The Law of Partition of Kinetic Energy". *Phil. Mag.*, **49**: 98–118.
28. Pais, A. (1982). "Subtle is the Lord". Oxford University Press. Oxford UK.
29. Armin, H. (1971). "The Genesis of Quantum Theory (1899–1913)". Translated by Claude W. Nash, ed.), Cambridge, MA.
30. Mayhew, K.W. (2017). "A New Perspective for Kinetic Theory and Heat Capacity". *Prog. in Phys.*, **13** (3): 165-173.
31. Liu, J.Z. (2023), "[The Nature of Absolute Zero Temperature](#)", Stanford University. Archived ([PDF](#)).
32. Liu, J.Z. (2023), "[Why Phase Transition Temperature Remains Constant](#)", Stanford University. Archived ([PDF](#)).