

Logic, Interaction and Collective Agency

Lecture 2

ESSLLI'10, Copenhagen

Eric Pacuit

Olivier Roy

TiLPS, Tilburg University
`ai.stanford.edu/~epacuit`

University of Groningen
`philos.rug.nl/~olivier`

August 18, 2010

Example (1)

Suppose there are two friends Ann and Bob are on a bus separated by a crowd.

Example (1)

Suppose there are two friends Ann and Bob are on a bus separated by a crowd. Before the bus comes to the next stop a mutual friend from outside the bus yells “get off at the next stop to get a drink?”.

Example (1)

Suppose there are two friends Ann and Bob are on a bus separated by a crowd. Before the bus comes to the next stop a mutual friend from outside the bus yells “get off at the next stop to get a drink?”.

Say Ann is standing near the front door and Bob near the back door.

Example (1)

Suppose there are two friends Ann and Bob are on a bus separated by a crowd. Before the bus comes to the next stop a mutual friend from outside the bus yells “get off at the next stop to get a drink?”.

Say Ann is standing near the front door and Bob near the back door. When the bus comes to a stop, will they get off?

Example (1)

Suppose there are two friends Ann and Bob are on a bus separated by a crowd. Before the bus comes to the next stop a mutual friend from outside the bus yells “get off at the next stop to get a drink?”.

Say Ann is standing near the front door and Bob near the back door. When the bus comes to a stop, will they get off?

D. Lewis. *Convention*. 1969.

M. Chwe. *Rational Ritual*. 2001.

Example (2)

A and B are players in the same football team. A has the ball, but an opposing player is converging on him.

Example (2)

A and *B* are players in the same football team. *A* has the ball, but an opposing player is converging on him. He can pass the ball to *B*, who has a chance to shoot.

Example (2)

A and B are players in the same football team. A has the ball, but an opposing player is converging on him. He can pass the ball to B , who has a chance to shoot. There are two directions in which A can move the ball, *left* and *right*, and correspondingly, two directions in which B can run to intercept the pass.

Example (2)

A and B are players in the same football team. A has the ball, but an opposing player is converging on him. He can pass the ball to B , who has a chance to shoot. There are two directions in which A can move the ball, *left* and *right*, and correspondingly, two directions in which B can run to intercept the pass. If both choose *left* there is a 10% chance that a goal will be scored.

Example (2)

A and B are players in the same football team. A has the ball, but an opposing player is converging on him. He can pass the ball to B , who has a chance to shoot. There are two directions in which A can move the ball, *left* and *right*, and correspondingly, two directions in which B can run to intercept the pass. If both choose *left* there is a 10% chance that a goal will be scored. If they both choose *right*, there is a 11% change.

Example (2)

A and B are players in the same football team. A has the ball, but an opposing player is converging on him. He can pass the ball to B , who has a chance to shoot. There are two directions in which A can move the ball, *left* and *right*, and correspondingly, two directions in which B can run to intercept the pass. If both choose *left* there is a 10% chance that a goal will be scored. If they both choose *right*, there is a 11% change. Otherwise, the chance is zero.

Example (2)

A and B are players in the same football team. A has the ball, but an opposing player is converging on him. He can pass the ball to B , who has a chance to shoot. There are two directions in which A can move the ball, *left* and *right*, and correspondingly, two directions in which B can run to intercept the pass. If both choose *left* there is a 10% chance that a goal will be scored. If they both choose *right*, there is a 11% change. Otherwise, the chance is zero. There is no time for communication; the two players must act simultaneously.

Example (2)

A and B are players in the same football team. A has the ball, but an opposing player is converging on him. He can pass the ball to B , who has a chance to shoot. There are two directions in which A can move the ball, *left* and *right*, and correspondingly, two directions in which B can run to intercept the pass. If both choose *left* there is a 10% chance that a goal will be scored. If they both choose *right*, there is a 11% change. Otherwise, the chance is zero. There is no time for communication; the two players must act simultaneously.

What should they do?

R. Sugden. *The Logic of Team Reasoning*. Philosophical Explorations (6)3, pgs. 165 - 181 (2003).

Example (2)

		<i>B</i>	
		<i>l</i>	<i>r</i>
<i>A</i>	<i>l</i>		
	<i>r</i>		

Example (2)

		<i>B</i>	
		<i>l</i>	<i>r</i>
<i>A</i>	<i>l</i>	10,10	0,0
	<i>r</i>	0,0	11,11

A: What should I do?

Example (2)

		<i>B</i>	
		<i>l</i>	<i>r</i>
<i>A</i>	<i>l</i>	10,10	0,0
	<i>r</i>	0,0	11,11

A: What should I do? *r* if the probability of *B* choosing *r* is $> \frac{10}{21}$
and *l* if the probability of *B* choosing *l* is $> \frac{11}{21}$
(symmetric reasoning for *B*)

Example (2)

		<i>B</i>	
		<i>l</i>	<i>r</i>
<i>A</i>	<i>l</i>	10,10	0,0
	<i>r</i>	0,0	11,11

A: What should I do? *r* if the probability of *B* choosing *r* is $> \frac{10}{21}$
and *l* if the probability of *B* choosing *l* is $> \frac{11}{21}$
(symmetric reasoning for *B*)

Example (2)

		<i>B</i>	
		<i>l</i>	<i>r</i>
<i>A</i>	<i>l</i>	10,10	0,0
	<i>r</i>	0,0	11,11

A: What should *we* do?

Example (2)

		B	
		l	r
A	l	10,10	0,0
	r	0,0	11,11

A: What should *we* do? **Team Reasoning**: an escape from the infinite regress? why should this “mode of reasoning” be endorsed?

Example (2)

		<i>B</i>	
		<i>l</i>	<i>r</i>
<i>A</i>	<i>l</i>	10,10	0,0
	<i>r</i>	0,0	11,11

A: What should *we* do? **Team Reasoning**: why should this “mode of reasoning” be endorsed?

Plan for Today

- ▶ Group Informational Attitudes

“*Common Knowledge*” is informally described as what any fool would know, given a certain situation: It encompasses what is relevant, agreed upon, established by precedent, assumed, being attended to, salient, or in the conversational record.

“*Common Knowledge*” is informally described as what any fool would know, given a certain situation: It encompasses what is relevant, agreed upon, established by precedent, assumed, being attended to, salient, or in the conversational record.

It is not Common Knowledge who “defined” Common Knowledge!

The first formal definition of common knowledge?

M. Friedell. *On the Structure of Shared Awareness*. Behavioral Science (1969).

R. Aumann. *Agreeing to Disagree*. Annals of Statistics (1976).

The first formal definition of common knowledge?

M. Friedell. *On the Structure of Shared Awareness*. Behavioral Science (1969).

R. Aumann. *Agreeing to Disagree*. Annals of Statistics (1976).

The first rigorous analysis of common knowledge

D. Lewis. *Convention, A Philosophical Study*. 1969.

The first formal definition of common knowledge?

M. Friedell. *On the Structure of Shared Awareness*. Behavioral Science (1969).

R. Aumann. *Agreeing to Disagree*. Annals of Statistics (1976).

The first rigorous analysis of common knowledge

D. Lewis. *Convention, A Philosophical Study*. 1969.

Fixed-point definition: $\gamma := i$ and j know that (φ and γ)

G. Harman. *Review of Linguistic Behavior*. Language (1977).

J. Barwise. *Three views of Common Knowledge*. TARK (1987).

The first formal definition of common knowledge?

M. Friedell. *On the Structure of Shared Awareness*. Behavioral Science (1969).

R. Aumann. *Agreeing to Disagree*. Annals of Statistics (1976).

The first rigorous analysis of common knowledge

D. Lewis. *Convention, A Philosophical Study*. 1969.

Fixed-point definition: $\gamma := i$ and j know that (φ and γ)

G. Harman. *Review of Linguistic Behavior*. Language (1977).

J. Barwise. *Three views of Common Knowledge*. TARK (1987).

Shared situation: There is a *shared situation* s such that (1) s entails φ , (2) s entails everyone knows φ , plus other conditions

H. Clark and C. Marshall. *Definite Reference and Mutual Knowledge*. 1981.

M. Gilbert. *On Social Facts*. Princeton University Press (1989).


P. Vanderschraaf and G. Sillari. "*Common Knowledge*", *The Stanford Encyclopedia of Philosophy* (2009).
<http://plato.stanford.edu/entries/common-knowledge/>.

The “Standard” Account

R. Aumann. *Agreeing to Disagree*. Annals of Statistics (1976).


R. Fagin, J. Halpern, Y. Moses and M. Vardi. *Reasoning about Knowledge*. MIT Press, 1995.

The “Standard” Account




W is a set of **states** or **worlds**.

The “Standard” Account




An **event/proposition** is any (definable) subset $E \subseteq W$

The “Standard” Account




At each state, agents are assigned a set of states they *consider possible* (according to their information).
The information may be (in)correct, partitional,

The “Standard” Account




Knowledge Function: $K_i : \wp(W) \rightarrow \wp(W)$ where
 $K_i(E) = \{w \mid R_i(w) \subseteq E\}$

The “Standard” Account




$$w \in K_A(E) \text{ and } w \notin K_B(E)$$

The “Standard” Account




The model also describes the agents' **higher-order knowledge/beliefs**

The “Standard” Account



Everyone Knows: $K(E) = \bigcap_{i \in \mathcal{A}} K_i(E)$, $K^0(E) = E$,
 $K^m(E) = K(K^{m-1}(E))$


The “Standard” Account



Common Knowledge: $C : \wp(W) \rightarrow \wp(W)$ with


$$C(E) = \bigcap_{m \geq 0} K^m(E)$$

The “Standard” Account



$$w \in K(E) \quad w \notin C(E)$$

The “Standard” Account



$$w \in C(E)$$

Fact. For all $i \in \mathcal{A}$ and $E \subseteq W$, $K_i C(E) = C(E)$.

Fact. For all $i \in \mathcal{A}$ and $E \subseteq W$, $K_i C(E) = C(E)$.

Suppose you are told “Ann and Bob are going together,” and respond “sure, that’s common knowledge.” What you mean is not only that everyone knows this, but also that the announcement is pointless, occasions no surprise, reveals nothing new; in effect, that the situation after the announcement does not differ from that before. ...the event “Ann and Bob are going together” — call it E — is common knowledge if and only if some event — call it F — happened that entails E and also entails all players’ knowing F (like all players met Ann and Bob at an intimate party). (*Aumann, pg. 271, footnote 8*)

Fact. For all $i \in \mathcal{A}$ and $E \subseteq W$, $K_i C(E) = C(E)$.

An event F is **self-evident** if $K_i(F) = F$ for all $i \in \mathcal{A}$.

Fact. An event E is commonly known iff some self-evident event that entails E obtains.

Fact. For all $i \in \mathcal{A}$ and $E \subseteq W$, $K_i C(E) = C(E)$.

An event F is **self-evident** if $K_i(F) = F$ for all $i \in \mathcal{A}$.

Fact. An event E is commonly known iff some self-evident event that entails E obtains.

Fact. $w \in C(E)$ if every finite path starting at w ends in a state in E

The following axiomatize common knowledge:

- ▶ $C(\varphi \rightarrow \psi) \rightarrow (C\varphi \rightarrow C\psi)$
- ▶ $C\varphi \rightarrow (\varphi \wedge EC\varphi)$ (Fixed-Point)
- ▶ $C(\varphi \rightarrow E\varphi) \rightarrow (\varphi \rightarrow C\varphi)$ (Induction)

An Example

Two players Ann and Bob are told that the following will happen. Some positive integer n will be chosen and *one* of n , $n + 1$ will be written on Ann's forehead, the other on Bob's. Each will be able to see the other's forehead, but not his/her own.

An Example

Two players Ann and Bob are told that the following will happen. Some positive integer n will be chosen and *one* of n , $n + 1$ will be written on Ann's forehead, the other on Bob's. Each will be able to see the other's forehead, but not his/her own.

Suppose the number are (2,3).

An Example

Two players Ann and Bob are told that the following will happen. Some positive integer n will be chosen and *one* of n , $n + 1$ will be written on Ann's forehead, the other on Bob's. Each will be able to see the other's forehead, but not his/her own.

Suppose the number are (2,3).

Do the agents know there numbers are less than 1000?


An Example

Two players Ann and Bob are told that the following will happen. Some positive integer n will be chosen and *one* of n , $n + 1$ will be written on Ann's forehead, the other on Bob's. Each will be able to see the other's forehead, but not his/her own.

Suppose the number are (2,3).

Do the agents know their numbers are less than 1000?

Is it common knowledge that their numbers are less than 1000?



The Fixed-Point Definition

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E))$

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E)) = K(E \cap C(E))$

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E)) = K(E \cap C(E)) = K(C(E)) = \bigcap_{i \in \mathcal{A}} K_i(C(E))$

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E)) = K(E \cap C(E)) = K(C(E)) = \bigcap_{i \in \mathcal{A}} K_i(C(E)) = \bigcap_{i \in \mathcal{A}} C(E) = C(E)$

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E)) = K(E \cap C(E)) = K(C(E)) = \bigcap_{i \in \mathcal{A}} K_i(C(E)) = \bigcap_{i \in \mathcal{A}} C(E) = C(E)$
- ▶ There are other fixed points of f_E : $f_E(\perp) = \perp$

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E)) = K(E \cap C(E)) = K(C(E)) = \bigcap_{i \in \mathcal{A}} K_i(C(E)) = \bigcap_{i \in \mathcal{A}} C(E) = C(E)$
- ▶ There are other fixed points of f_E : $f_E(\perp) = \perp$
- ▶ f_E is monotonic:

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E)) = K(E \cap C(E)) = K(C(E)) = \bigcap_{i \in \mathcal{A}} K_i(C(E)) = \bigcap_{i \in \mathcal{A}} C(E) = C(E)$
- ▶ There are other fixed points of f_E : $f_E(\perp) = \perp$
- ▶ f_E is monotonic: $A \subseteq B$ implies $E \cap A \subseteq E \cap B$.

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E)) = K(E \cap C(E)) = K(C(E)) = \bigcap_{i \in \mathcal{A}} K_i(C(E)) = \bigcap_{i \in \mathcal{A}} C(E) = C(E)$
- ▶ There are other fixed points of f_E : $f_E(\perp) = \perp$
- ▶ f_E is monotonic: $A \subseteq B$ implies $E \cap A \subseteq E \cap B$. Then $f_E(E \cap A) = K(E \cap A) \subseteq K(E \cap B) = f_E(E \cap B)$

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E)) = K(E \cap C(E)) = K(C(E)) = \bigcap_{i \in \mathcal{A}} K_i(C(E)) = \bigcap_{i \in \mathcal{A}} C(E) = C(E)$
- ▶ There are other fixed points of f_E : $f_E(\perp) = \perp$
- ▶ f_E is monotonic: $A \subseteq B$ implies $E \cap A \subseteq E \cap B$. Then $f_E(E \cap A) = K(E \cap A) \subseteq K(E \cap B) = f_E(E \cap B)$
- ▶ (Tarski) Every monotone operator has a greatest (and least) fixed point

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E)) = K(E \cap C(E)) = K(C(E)) = \bigcap_{i \in \mathcal{A}} K_i(C(E)) = \bigcap_{i \in \mathcal{A}} C(E) = C(E)$
- ▶ There are other fixed points of f_E : $f_E(\perp) = \perp$
- ▶ f_E is monotonic: $A \subseteq B$ implies $E \cap A \subseteq E \cap B$. Then $f_E(E \cap A) = K(E \cap A) \subseteq K(E \cap B) = f_E(E \cap B)$
- ▶ (Tarski) Every monotone operator has a greatest (and least) fixed point
- ▶ Let $K^*(E)$ be the greatest fixed point of f_E .

The Fixed-Point Definition

$$f_E(X) = K(E \cap X) = \bigcap_{i \in \mathcal{A}} K_i(E \cap X)$$

- ▶ $C(E)$ is a fixed point of f_E : $f_E(C(E)) = K(E \cap C(E)) = K(C(E)) = \bigcap_{i \in \mathcal{A}} K_i(C(E)) = \bigcap_{i \in \mathcal{A}} C(E) = C(E)$
- ▶ There are other fixed points of f_E : $f_E(\perp) = \perp$
- ▶ f_E is monotonic: $A \subseteq B$ implies $E \cap A \subseteq E \cap B$. Then $f_E(E \cap A) = K(E \cap A) \subseteq K(E \cap B) = f_E(E \cap B)$
- ▶ (Tarski) Every monotone operator has a greatest (and least) fixed point
- ▶ Let $K^*(E)$ be the greatest fixed point of f_E .
- ▶ **Fact.** $K^*(E) = C(E)$.

The Fixed-Point Definition

Separating the fixed-point/iteration definition of common knowledge/belief:

J. Barwise. *Three views of Common Knowledge*. TARK (1987).

J. van Benthem and D. Saraenac. *The Geometry of Knowledge*. Aspects of Universal Logic (2004).

A. Heifetz. *Iterative and Fixed Point Common Belief*. Journal of Philosophical Logic (1999).

Some Issues

Some Issues

- ▶ What *does* a group know/believe/accept? vs. what *can* a group (come to) know/believe/accept?

Some Issues

- ▶ What *does* a group know/believe/accept? vs. what *can* a group (come to) know/believe/accept?

C. List. *Group knowledge and group rationality: a judgment aggregation perspective*. Episteme (2008).

Some Issues

- ▶ What *does* a group know/believe/accept? vs. what *can* a group (come to) know/believe/accept?

C. List. *Group knowledge and group rationality: a judgment aggregation perspective*. Episteme (2008).

- ▶ Other “group informational attitudes”: distributed knowledge, common belief, ...

Some Issues

- ▶ What *does* a group know/believe/accept? vs. what *can* a group (come to) know/believe/accept?

C. List. *Group knowledge and group rationality: a judgment aggregation perspective*. Episteme (2008).

- ▶ Other “group informational attitudes”: distributed knowledge, common belief, ...
- ▶ Levels of knowledge

Some Issues

- ▶ What *does* a group know/believe/accept? vs. what *can* a group (come to) know/believe/accept?

C. List. *Group knowledge and group rationality: a judgment aggregation perspective*. Episteme (2008).

- ▶ Other “group informational attitudes”: distributed knowledge, common belief, ...
- ▶ Levels of knowledge
- ▶ Common knowledge/belief of *rationality*

Some Issues

- ▶ What *does* a group know/believe/accept? vs. what *can* a group (come to) know/believe/accept?

C. List. *Group knowledge and group rationality: a judgment aggregation perspective*. Episteme (2008).

- ▶ Other “group informational attitudes”: distributed knowledge, common belief, ...
- ▶ Levels of knowledge
- ▶ Common knowledge/belief of *rationality*
- ▶ Where does common knowledge come from?

Some Issues

- ✓ What *does* a group know/believe/accept? vs. what *can* a group (come to) know/believe/accept?

C. List. *Group knowledge and group rationality: a judgment aggregation perspective*. Episteme (2008).

- ▶ Other “group informational attitudes”: distributed knowledge, common belief, ...
- ▶ Levels of knowledge
- ▶ Common knowledge/belief of *rationality*
- ▶ Where does common knowledge come from?

Distributed Knowledge

$$D_G(E) = \{w \mid \left(\bigcap_{i \in G} R_i(w) \right) \subseteq E\}$$

Distributed Knowledge

$$D_G(E) = \{w \mid \left(\bigcap_{i \in G} R_i(w) \right) \subseteq E\}$$

- ▶ $K_A(p) \wedge K_B(p \rightarrow q) \rightarrow D_{A,B}(q)$
- ▶ $D_G(\varphi) \rightarrow \bigwedge_{i \in G} K_i \varphi$

Distributed Knowledge

$$D_G(E) = \{w \mid \left(\bigcap_{i \in G} R_i(w) \right) \subseteq E\}$$

- ▶ $K_A(p) \wedge K_B(p \rightarrow q) \rightarrow D_{A,B}(q)$
- ▶ $D_G(\varphi) \rightarrow \bigwedge_{i \in G} K_i \varphi$

F. Roelofsen. *Distributed Knowledge*. Journal of Applied Nonclassical Logic (2006).

Distributed Knowledge

$$D_G(E) = \{w \mid \left(\bigcap_{i \in G} R_i(w) \right) \subseteq E\}$$

- ▶ $K_A(p) \wedge K_B(p \rightarrow q) \rightarrow D_{A,B}(q)$
- ▶ $D_G(\varphi) \rightarrow \bigwedge_{i \in G} K_i \varphi$

F. Roelofsen. *Distributed Knowledge*. Journal of Applied Nonclassical Logic (2006).

$w \in K_G(E)$ iff $R_G(w) \subseteq E$ (without necessarily $R_G(w) = \bigcap_{i \in G} R_i(w)$)

A. Baltag and S. Smets. *Correlated Knowledge: an Epistemic-Logic view on Quantum Entanglement*. Int. Journal of Theoretical Physics (2010).

Common p -belief

The typical example of an event that creates common knowledge is a **public announcement**.

Common p -belief

The typical example of an event that creates common knowledge is a **public announcement**.

Shouldn't one always allow for some small probability that a participant was absentminded, not listening, sending a text, checking facebook, proving a theorem, asleep, ...

Common p -belief

The typical example of an event that creates common knowledge is a **public announcement**.

Shouldn't one always allow for some small probability that a participant was absentminded, not listening, sending a text, checking facebook, proving a theorem, asleep, ...

“We show that the weaker concept of “common belief” can function successfully as a substitute for common knowledge in the theory of equilibrium of Bayesian games.”

D. Monderer and D. Samet. *Approximating Common Knowledge with Common Beliefs*. Games and Economic Behavior (1989).

Common p -belief: definition

Let $(W, \{R_i\}_{i \in \mathcal{A}}, p, V)$ be a Bayesian model.

Common p -belief: definition

Let $(W, \{R_i\}_{i \in \mathcal{A}}, p, V)$ be a Bayesian model.

$$B_i^p(E) = \{w \mid p(E \mid R_i(w)) \geq p\}$$

Common p -belief: definition

Let $(W, \{R_i\}_{i \in \mathcal{A}}, p, V)$ be a Bayesian model.

$$B_i^p(E) = \{w \mid p(E \mid R_i(w)) \geq p\}$$

An event E is **evident p -belief** if for each $i \in \mathcal{A}$, $E \subseteq B_i^p(E)$

Common p -belief: definition


Let $(W, \{R_i\}_{i \in \mathcal{A}}, p, V)$ be a Bayesian model.

$$B_i^p(E) = \{w \mid p(E \mid R_i(w)) \geq p\}$$

An event E is **evident p -belief** if for each $i \in \mathcal{A}$, $E \subseteq B_i^p(E)$


An event F is **common p -belief** at w if there exists an evident p -belief event E such that $w \in E$ and for all $i \in \mathcal{A}$, $E \subseteq B_i^p(F)$

Common p -belief: example




Two agents either hear (H) or don't hear (D) the announcement.

Common p -belief: example




The probability that an agent hears is $1 - \epsilon$.

Common p -belief: example




The agents *know* their “type”.

Common p -belief: example




The event “everyone hears” ($E = \{w_1\}$)

Common p -belief: example




The event “everyone hears” ($E = \{w_1\}$) is **not** common knowledge

Common p -belief: example



The event “everyone hears” ($E = \{w_1\}$) is **not** common knowledge, but it is **common $(1 - \epsilon)$ -belief**

Common p -belief: example



The event “everyone hears” ($E = \{w_1\}$) is **not** common knowledge, but it is **common $(1 - \epsilon)$ -belief**:

$B_i^{(1-\epsilon)}(E) = \{w \mid p(E \mid \Pi_i(w)) \geq 1 - \epsilon\} = \{w_1\} = E$,
for $i = 1, 2$

Some Issues

- ✓ What *does* a group know/believe/accept? vs. what *can* a group (come to) know/believe/accept?

C. List. *Group knowledge and group rationality: a judgment aggregation perspective*. Episteme (2008).

- ✓ Other “group informational attitudes”: distributed knowledge, common belief, ...
 - ▶ Levels of knowledge
 - ▶ Common knowledge/belief of *rationality*
 - ▶ Where does common knowledge come from?

Levels of Knowledge

Levels of Knowledge

What are the *states of knowledge* created in a group when communication takes place? What happens when communication is not the the whole group, but pairwise?

R. Parikh and P. Krasucki. *Communication, Consensus and Knowledge*. Journal of Economic Theory (1990).

Informal Definition: Given some fact P and a set of agents \mathcal{A} , a **state of knowledge** is a (consistent) description of the agents first-order and higher-order information about P .

Levels of Knowledge

What are the *states of knowledge* created in a group when communication takes place? What happens when communication is not the the whole group, but pairwise?

R. Parikh and P. Krasucki. *Communication, Consensus and Knowledge*. Journal of Economic Theory (1990).

Informal Definition: Given some fact P and a set of agents \mathcal{A} , a **state of knowledge** is a (consistent) description of the agents first-order and higher-order information about P .

For example: $\{P, K_A P, K_B P, K_B K_A P\}$

Levels of Knowledge

Informal Definition: Given some fact P and a set of agents \mathcal{A} , a **state of knowledge** is a (consistent) description of the agents first-order and higher-order information about P .

For example: $\{P, K_A P, K_B P, K_B K_A P\}$

Also called **level of knowledge**, **hierarchy of knowledge**.

Levels of Knowledge

Informal Definition: Given some fact P and a set of agents \mathcal{A} , a **state of knowledge** is a (consistent) description of the agents first-order and higher-order information about P .

For example: $\{P, K_A P, K_B P, K_B K_A P\}$

Also called **level of knowledge**, **hierarchy of knowledge**.

At one extreme, no one may have any information about P and the other extreme is when there is common knowledge of P .

There are many interesting levels in between...

Levels of Knowledge

Informal Definition: Given some fact P and a set of agents \mathcal{A} , a **state of knowledge** is a (consistent) description of the agents first-order and higher-order information about P .

For example: $\{P, K_i P, K_j P, K_j K_i P\}$

Also called **level of knowledge**, **hierarchy of knowledge**.

At one extreme, no one may have any information about P and the other extreme is when there is **common knowledge** of P .

There are many interesting levels in between...

Levels of Knowledge

Informal Definition: Given some fact P and a set of agents \mathcal{A} , a **state of belief** is a (consistent) description of the agents first-order and higher-order information about P .

For example: $\{P, B_i P, B_j P, B_j B_i P\}$

Also called **level of belief**, **hierarchy of belief**.

At one extreme, no one may have any information about P and the other extreme is when there is **common belief** of P .

There are many interesting levels in between...

Some Questions/Issues

- ▶ How do states of knowledge influence decisions in *game situations*?
- ▶ Can we *realize* any state of knowledge?
- ▶ What is a *state* in an epistemic model?
- ▶ Is an epistemic model *common knowledge* among the agents?

States of Knowledge in Games

R. Parikh. *Levels of knowledge, games and group action*. Research in Economics 57, pp. 267 - 281 (2003).

States of Knowledge in Games

	G	N
g		
n		

States of Knowledge in Games

	G	N
g		$(1, 0)$
n	$(0, 1)$	

States of Knowledge in Games

	G	N
g		$(1, 0)$
n	$(0, 1)$	$(0, 0)$

States of Knowledge in Games

	G	N
g	$(-100, -10)$	$(1, 0)$
n	$(0, 1)$	$(0, 0)$

States of Knowledge in Games

	G	N
g	$(-100, -10)$	$(1, 0)$
n	$(0, 1)$	$(0, 0)$

States of Knowledge in Games

	G	N
g	$(-100, -10)$	$(1, 0)$
n	$(0, 1)$	$(0, 0)$

States of Knowledge in Games

	G	N
g	$(-100, -10)$	$(1, 0)$
n	$(0, 1)$	$(0, 0)$

$C_{p,mC}$

States of Knowledge in Games

	G	N
g	$(-100, -10)$	$(1, 0)$
n	$(0, 1)$	$(0, 0)$

$$K_p c, \neg K_m K_p c$$

Realizing States of Knowledge

What is the exact relationship between

Realizing States of Knowledge

What is the exact relationship between 1. types of communicatory events (public broadcasts, private announcements, DEL events, etc.),

Realizing States of Knowledge

What is the exact relationship between 1. types of communicatory events (public broadcasts, private announcements, DEL events, etc.), 2. the *protocol* (eg., synchronous, asynchronous, communication graph) and

Realizing States of Knowledge

What is the exact relationship between 1. types of communicatory events (public broadcasts, private announcements, DEL events, etc.), 2. the *protocol* (eg., synchronous, asynchronous, communication graph) and 3. different levels of knowledge?

Realizing States of Knowledge

What is the exact relationship between 1. types of communicatory events (public broadcasts, private announcements, DEL events, etc.), 2. the *protocol* (eg., synchronous, asynchronous, communication graph) and 3. different levels of knowledge?

Classic example: general's problem or the email problem show that common knowledge cannot be realized in systems with only asynchronous communication.

(cf. Halpern and Moses, Rubinstein)

Realizing States of Knowledge

What is the exact relationship between 1. types of communicatory events (public broadcasts, private announcements, DEL events, etc.), 2. the *protocol* (eg., synchronous, asynchronous, communication graph) and 3. different levels of knowledge?

Classic example: general's problem or the email problem show that common knowledge cannot be realized in systems with only asynchronous communication.

(cf. Halpern and Moses, Rubinstein)

What about other levels of knowledge?

R. Parikh and P. Krasucki. *Levels of knowledge in distributed computing*. Sadhana-Proceedings of the Indian Academy of Science 17 (1992).

What is a *State*?

Possible worlds, or states, are taken as primitive in Kripke structures. But in many applications, we intuitively understand what a state *is*:

What is a *State*?

Possible worlds, or states, are taken as primitive in Kripke structures. But in many applications, we intuitively understand what a state *is*:

Dynamic logic: a program state (assignment of values to variables)

Temporal logic: a moment in time

Distributed system: a sequence of local states for each process

What is a *State*?

Possible worlds, or states, are taken as primitive in Kripke structures. But in many applications, we intuitively understand what a state *is*:

Dynamic logic: a program state (assignment of values to variables)

Temporal logic: a moment in time

Distributed system: a sequence of local states for each process

What about in *game situations*?

What is a *State*?

Possible worlds, or states, are taken as primitive in Kripke structures. But in many applications, we intuitively understand what a state *is*:

Dynamic logic: a program state (assignment of values to variables)

Temporal logic: a moment in time

Distributed system: a sequence of local states for each process

What about in *game situations*?

Answer: a *description* of the first-order and higher-order information of the players

R. Fagin, J. Halpern and M. Vardi. *Model theoretic analysis of knowledge*. Journal of the ACM 91 (1991).

Is an Epistemic Model “Common Knowledge”?

“The implicit assumption that the information partitions...are themselves common knowledge...constitutes no loss of generality... the assertion that each individual ‘knows’ the knowledge operators of all individual has no real substance; it is part of the framework.”

R. Aumann. *Interactive Epistemology I & II*. International Journal of Game Theory (1999).

Is an Epistemic Model “Common Knowledge” ?

“The implicit assumption that the information partitions...are themselves common knowledge...constitutes no loss of generality... the assertion that each individual ‘knows’ the knowledge operators of all individual has no real substance; it is part of the framework.”

R. Aumann. *Interactive Epistemology I & II*. International Journal of Game Theory (1999).

“it is an informal but *meaningful* meta-assumption....It is not trivial at all to assume it is “common knowledge” which partition every player has.”

A. Heifetz. *How canonical is the canonical model? A comment on Aumann's interactive epistemology*. International Journal of Game Theory (1999).

A General Question

How many levels/states of knowledge (beliefs) are there?

A General Question

How many levels/states of knowledge (beliefs) are there?

It depends on how you count:

- ▶ Parikh and Krasucki: Countably many *levels* of knowledge

▶ Why?

- ▶ Parikh and EP: Uncountably many levels of belief

▶ Why?

- ▶ Hart, Heiftetz and Samet: Uncountably many *states* of knowledge

▶ Why?

Returning to the Motivating Questions

- ▶ How do states of knowledge influence decisions in *game situations*?
- ▶ Can we *realize* any state of knowledge?
- ▶ What is a *state* in an epistemic model?
- ▶ Is an epistemic model *common knowledge* among the agents?

Returning to the Motivating Questions

- ▶ How do states of knowledge influence decisions in *game situations*?
- ▶ Can we *realize* any state of knowledge?
- ▶ What is a *state* in an epistemic model?
- ▶ Is an epistemic model *common knowledge* among the agents?

Returning to the Motivating Questions

- ▶ How do states of knowledge influence decisions in *game situations*?
- ▶ Can we *realize* any state of knowledge?
It depends...
- ▶ What is a *state* in an epistemic model?
It depends...
- ▶ Is an epistemic model *common knowledge* among the agents?
It depends...


Some Issues

- ✓ What *does* a group know/believe/accept? vs. what *can* a group (come to) know/believe/accept?


C. List. *Group knowledge and group rationality: a judgment aggregation perspective*. Episteme (2008).

- ✓ the “group informational attitudes”: distributed knowledge, common belief, ...
- ✓ Levels of knowledge
 - ▶ Common knowledge/belief of *rationality*
 - ▶ Where does common knowledge come from?


“Rational” Strategies



“Rational” Strategies




“Rational” Strategies



- Q1: Can we always find a model where $Rat \neq \emptyset$?

“Rational” Strategies



- ▶ Q1: Can we always find a model where $Rat \neq \emptyset$?
- ▶ Q2: Can we *characterize* the strategies that are *always* in Rat ?

Some Properties

- ▶ Knowledge of own choice
the agents choices are uniform in their information sets

Some Properties

- ▶ Knowledge of own choice
the agents choices are uniform in their information sets
- ▶ The structure of the game is commonly known
the map from states to strategy profiles is onto

Some Properties

- ▶ Knowledge of own choice
the agents choices are uniform in their information sets
- ▶ The structure of the game is commonly known
the map from states to strategy profiles is onto
- ▶ Common prior
all agents have the same prior probability function

Some Properties

- ▶ Knowledge of own choice
the agents choices are uniform in their information sets
- ▶ The structure of the game is commonly known
the map from states to strategy profiles is onto
- ▶ Common prior
all agents have the same prior probability function

S. Morris. *The common prior assumption in economic theory*. Economics and Philosophy, 11, pgs. 227 - 254, 1995.

Some Properties

- ▶ Knowledge of own choice
the agents choices are uniform in their information sets
- ▶ The structure of the game is commonly known
the map from states to strategy profiles is onto
- ▶ Common prior
all agents have the same prior probability function

S. Morris. *The common prior assumption in economic theory*. Economics and Philosophy, 11, pgs. 227 - 254, 1995.

- ▶ *Common Knowledge* of “rational choice”
there is no “Ann-Bob path” that leads outside of Rat

Returning to the questions

Simple Characterization Theorem In any *Bayesian model* of a finite strategic game, (the projection of) any state where the players are *rational* and there is *common knowledge of rationality* is exactly the set of strategies that survive *iterated removal of strictly dominated strategies*. (Question 2)

A. Brandenburger and E. Dekel. *Rationalizability and correlated equilibria*. *Econometrica*, 55:1391-1402, 1987.

Common Knowledge of Rationality

		Bob	
		<i>L</i>	<i>R</i>
Ann	<i>U</i>	1,2	0,1
	<i>D</i>	0,1	1,0

Common Knowledge of Rationality

		Bob	
		<i>L</i>	<i>R</i>
Ann	<i>U</i>	1,2	0,1
	<i>D</i>	0,1	1,0

There is no prior such that R is rational for Bob.

Common Knowledge of Rationality

		Bob	
		<i>L</i>	<i>R</i>
Ann	<i>U</i>	1,2	0,1
	<i>D</i>	0,1	1,0

If Ann knows this, then she does not consider *R* a option for Bob

Common Knowledge of Rationality

		Bob	
		<i>L</i>	<i>R</i>
Ann	<i>U</i>	1,2	0,1
	<i>D</i>	0,1	1,0

So, *U* is the only rational choice.

Other natural properties...

- ▶ Do not *initially* rule out any strategies of the other players (admissibility)
- ▶ If two strategies are rational for an opponent, then neither can be “ruled out” (picking vs. choosing: *i* knows which options *j* will choose from, but *i* cannot know which optioned *j* picked)
- ▶ Do not *initially* rule out any *types* of the other players

...lead to puzzles

R. Cubitt and R. Sugden. *Rationally Justifiable Play and the Theory of Non-cooperative games*. Economic Journal, 104, pgs. 798 - 803, 1994.

R. Cubitt and R. Sugden. *Common reasoning in games*. Manuscript, 2008.

A. Brandenburger and H. J. Keisler. *An Impossibility Theorem on Beliefs in Games*. *Studia Logica* (2006).

Proving too Much Puzzle

		Bob	
		L	R
Ann	F	0,0	0,0
	S	-1,3	2,2
	T	-1,3	1,5

Proposition In every* Bayesian model satisfying *privacy of tie-breaking* of the above game, $Rat_A = \{F\}$ & $Rat_B = \{L, R\}$

Then player 1 must assign probability greater than $\frac{2}{3}$ to player 2 playing L. But why is this justified?


Proving too Much Puzzle

		Bob	
		L	R
Ann	F	0,0	0,0
	S	-1,3	2,2
	T	-1,3	1,5

Proposition In every* Bayesian model satisfying *privacy of tie-breaking* of the above game, $Rat_A = \{F\}$ & $Rat_B = \{L, R\}$


Then player 1 must assign probability greater than $\frac{2}{3}$ to player 2 playing L. But why is this justified?

		Bob	
		<i>L</i>	<i>R</i>
Ann	<i>F</i>	0,0	0,0
	<i>S</i>	-1,3	2,2
	<i>T</i>	-1,3	1,5




		Bob	
		<i>L</i>	<i>R</i>
Ann	<i>F</i>	0,0	0,0
	<i>S</i>	-1,3	2,2
	<i>T</i>	-1,3	1,5


1. *T* cannot be rational for Ann




		Bob	
		<i>L</i>	<i>R</i>
Ann	<i>F</i>	0,0	0,0
	<i>S</i>	-1,3	2,2
	<i>T</i>	-1,3	1,5



1. *T* cannot be rational for Ann
2. if Bob assigns a nonzero probability to *S*, then *R* is not rational




1. *T* cannot be rational for Ann
2. if Bob assigns a nonzero probability to *S*, then *R* is not rational
3. *S* is rational for Ann provided she considers it possible that Bob plays *R*




1. T cannot be rational for Ann
2. if Bob assigns a nonzero probability to S , then R is not rational
3. S is rational for Ann provided she considers it possible that Bob plays R

		Bob	
		<i>L</i>	<i>R</i>
Ann	<i>F</i>	0,0	0,0
	<i>S</i>	-1,3	2,2
	<i>T</i>	-1,3	1,5

1. *T* cannot be rational for Ann
2. if Bob assigns a nonzero probability to *S*, then *R* is not rational
3. *S* is rational for Ann provided she considers it possible that Bob plays *R*
4. *F* is the only rational choice for Ann



		Bob	
		<i>L</i>	<i>R</i>
Ann	<i>F</i>	0,0	0,0
	<i>S</i>	-1,3	2,2
	<i>T</i>	-1,3	1,5



1. *T* cannot be rational for Ann
2. if Bob assigns a nonzero probability to *S*, then *R* is not rational
3. *S* is rational for Ann provided she considers it possible that Bob plays *R*
4. *F* is the only rational choice for Ann
5. both *L* and *R* are rational responses if it is commonly known that Ann will play *F*

Paradox

Paradox

	in_2	out_2
in_1	1, 1, 1	1, 1, 1
out_1	1, 1, 1	0, 1, 1
	in_3	

	in_2	out_2
in_1	1, 1, 1	1, 0, 1
out_1	1, 1, 0	0, 0, 0
	out_3	

There is no Bayesian model of the above game satisfying privacy of tie-breaking.

Paradox

	in_2	out_2
in_1	1, 1, 1	1, 1, 1
out_1	1, 1, 1	0, 1, 1

in_3

	in_2	out_2
in_1	1, 1, 1	1, 0, 1
out_1	1, 1, 0	0, 0, 0

out_3

1. If 1 considers out_2 possible, then it is common knowledge that out_1 is not possible

Paradox

	in_2	out_2
in_1	1, 1, 1	1, 1, 1
out_1	1, 1, 1	0, 1, 1
	in_3	

	in_2	out_2
in_1	1, 1, 1	1, 0, 1
out_1	1, 1, 0	0, 0, 0
	out_3	

1. If 1 considers out_2 possible, then it is common knowledge that out_1 is not possible
2. If 2 considers out_3 possible, then it is common knowledge that out_2 is not possible

Paradox

	in_2	out_2
in_1	1, 1, 1	1, 1, 1
out_1	1, 1, 1	0, 1, 1
	in_3	

	in_2	out_2
in_1	1, 1, 1	1, 0, 1
out_1	1, 1, 0	0, 0, 0
	out_3	

1. If 1 considers out_2 possible, then it is common knowledge that out_1 is not possible
2. If 2 considers out_3 possible, then it is common knowledge that out_2 is not possible
3. If 3 considers out_1 possible, then it is common knowledge that out_3 is not possible

Paradox

	<i>in</i> ₂	<i>out</i> ₂
<i>in</i> ₁	1, 1, 1	1, 1, 1
<i>out</i> ₁	1, 1, 1	0, 1, 1
	<i>in</i> ₃	

	<i>in</i> ₂	<i>out</i> ₂
<i>in</i> ₁	1, 1, 1	1, 0, 1
<i>out</i> ₁	1, 1, 0	0, 0, 0
	<i>out</i> ₃	

4. If 1 does not consider *out*₂ possible, then 2 & 3 must consider *in*₁ & *out*₁ possible

Paradox

	<i>in</i> ₂	<i>out</i> ₂
<i>in</i> ₁	1, 1, 1	1, 1, 1
<i>out</i> ₁	1, 1, 1	0, 1, 1
	<i>in</i> ₃	

	<i>in</i> ₂	<i>out</i> ₂
<i>in</i> ₁	1, 1, 1	1, 0, 1
<i>out</i> ₁	1, 1, 0	0, 0, 0
	<i>out</i> ₃	

4. If 1 does not consider *out*₂ possible, then 2 & 3 must consider *in*₁ & *out*₁ possible
5. If 2 does not consider *out*₃ possible, then 1 & 3 must consider *in*₂ & *out*₂ possible

Paradox

	in_2	out_2
in_1	1, 1, 1	1, 1, 1
out_1	1, 1, 1	0, 1, 1
	in_3	

	in_2	out_2
in_1	1, 1, 1	1, 0, 1
out_1	1, 1, 0	0, 0, 0
	out_3	

4. If 1 does not consider out_2 possible, then 2 & 3 must consider in_1 & out_1 possible
5. If 2 does not consider out_3 possible, then 1 & 3 must consider in_2 & out_2 possible
6. If 3 does not consider out_1 possible, then 1 & 2 must consider in_3 & out_3 possible

Paradox

	in_2	out_2
in_1	1, 1, 1	1, 1, 1
out_1	1, 1, 1	0, 1, 1
	in_3	

	in_2	out_2
in_1	1, 1, 1	1, 0, 1
out_1	1, 1, 0	0, 0, 0
	out_3	

- ▶ If i considers out_{i+1} possible, then it is common knowledge that out_i is not possible
- ▶ If i does not consider out_{i+1} possible, then $i + 1$ & $i + 2$ must consider in_i & out_i possible

Paradox

	in_2	out_2
in_1	1, 1, 1	1, 1, 1
out_1	1, 1, 1	0, 1, 1
	in_3	

	in_2	out_2
in_1	1, 1, 1	1, 0, 1
out_1	1, 1, 0	0, 0, 0
	out_3	

- ▶ If i considers out_{i+1} possible, then it is common knowledge that out_i is not possible
- ▶ If i does not consider out_{i+1} possible, then $i + 1$ & $i + 2$ must consider in_i & out_i possible
- ▶ 1 **does consider** out_2 possible \implies 3 does not consider out_1 possible \implies 2 considers out_3 possible \implies 1 **does not consider** out_2 possible

Digression: Admissibility

Related to an old puzzle in game theory about admissibility (best response to some belief of player i with positive probability on all the strategies of the opponents):

Digression: Admissibility

Related to an old puzzle in game theory about admissibility (best response to some belief of player i with positive probability on all the strategies of the opponents):

A strategy is admissibility iff it is not *weakly dominated*.

D. Pearce. *Rationalizable strategic behavior and the problem of perfection*.
Econometrica, 52, pgs. 1029 - 1050, 1984.

Digression: Admissibility

Related to an old puzzle in game theory about admissibility (best response to some belief of player i with positive probability on all the strategies of the opponents):

A strategy is admissibility iff it is not *weakly dominated*.

D. Pearce. *Rationalizable strategic behavior and the problem of perfection*.
Econometrica, 52, pgs. 1029 - 1050, 1984.

Does assuming that it is commonly known that players play only admissible strategies lead to a process of iterated removal of weakly dominated strategies?

Digression: Admissibility

Related to an old puzzle in game theory about admissibility (best response to some belief of player i with positive probability on all the strategies of the opponents):

A strategy is admissibility iff it is not *weakly dominated*.

D. Pearce. *Rationalizable strategic behavior and the problem of perfection*. *Econometrica*, 52, pgs. 1029 - 1050, 1984.

Does assuming that it is commonly known that players play only admissible strategies lead to a process of iterated removal of weakly dominated strategies? **no!**

L. Samuelson. *Dominated Strategies and Common Knowledge*. *Games and Economic Behavior* (1992).

► Explanation

Some Issues

- ✓ What *does* a group know/believe/accept? vs. what *can* a group (come to) know/believe/accept?

C. List. *Group knowledge and group rationality: a judgment aggregation perspective*. Episteme (2008).

- ✓ Other “group informational attitudes”: distributed knowledge, common belief, ...
- ✓ Levels of knowledge
- ✓ Common knowledge/belief of *rationality*
- ▶ Where does common knowledge come from?

R. Cubitt and R. Sugden. *Common Knowledge, Salience and Convention: A Reconstruction of David Lewis' Game Theory*. *Economics and Philosophy*, 19, pgs. 175-210 , 2003..

Reason to Believe

$B_i\varphi$: “*i* believes φ ”

Reason to Believe

$B_i\varphi$: “*i* believes φ ” vs. $R_i(\varphi)$: “*i* has a *reason to believe* φ ”

Reason to Believe

$B_i\varphi$: “*i* believes φ ” vs. $R_i(\varphi)$: “*i* has a *reason to believe* φ ”

- ▶ “Although it is an essential part of Lewis’ theory that human beings are *to some degree* rational, he does not want to make the strong rationality assumptions of conventional decision theory or game theory.” (CS, pg. 184).

Reason to Believe

$B_i\varphi$: “*i* believes φ ” vs. $R_i(\varphi)$: “*i* has a *reason to believe* φ ”

- ▶ “Although it is an essential part of Lewis’ theory that human beings are *to some degree* rational, he does not want to make the strong rationality assumptions of conventional decision theory or game theory.” (CS, pg. 184).
- ▶ Anyone who accept the rules of arithmetic has a reason to believe $618 \times 377 = 232,986$, but most of us do not hold have firm beliefs about this.

Reason to Believe

$B_i\varphi$: “ i believes φ ” vs. $R_i(\varphi)$: “ i has a *reason to believe* φ ”

- ▶ “Although it is an essential part of Lewis’ theory that human beings are *to some degree* rational, he does not want to make the strong rationality assumptions of conventional decision theory or game theory.” (CS, pg. 184).
- ▶ Anyone who accept the rules of arithmetic has a reason to believe $618 \times 377 = 232,986$, but most of us do not hold have firm beliefs about this.
- ▶ Definition: $R_i(\varphi)$ means φ is true within some logic of reasoning that is *endorsed* by (that is, accepted as a normative standard by) person i ... φ must be either regarded as *self-evident* or derivable by rules of inference (deductive or inductive)

A indicates to i that φ

A is a “state of affairs”

$A \text{ ind}_i \varphi$: i 's reason to believe that A holds *provides* i 's reason for believing that φ is true.

(A1) For all i , for all A , for all φ : $[R_i(A \text{ holds}) \wedge (A \text{ ind}_i \varphi)] \Rightarrow R_i(\varphi)$

Some Properties

Some Properties

- ▶ $[(A \text{ holds}) \text{ entails } (A' \text{ holds})] \Rightarrow A \text{ ind}_i(A' \text{ holds})$

Some Properties

- ▶ $[(A \text{ holds}) \text{ entails } (A' \text{ holds})] \Rightarrow A \text{ ind}_i(A' \text{ holds})$
- ▶ $[(A \text{ ind}_i \varphi) \wedge (A \text{ ind}_i \psi)] \Rightarrow A \text{ ind}_i(\varphi \wedge \psi)$

Some Properties

- ▶ $[(A \text{ holds}) \text{ entails } (A' \text{ holds})] \Rightarrow A \text{ ind}_i(A' \text{ holds})$
- ▶ $[(A \text{ ind}_i \varphi) \wedge (A \text{ ind}_i \psi)] \Rightarrow A \text{ ind}_i(\varphi \wedge \psi)$
- ▶ $[(A \text{ ind}_i[A' \text{ holds}]) \wedge (A' \text{ ind}_i x)] \Rightarrow A \text{ ind}_i \varphi$

Some Properties

- ▶ $[(A \text{ holds}) \text{ entails } (A' \text{ holds})] \Rightarrow A \text{ ind}_i(A' \text{ holds})$
- ▶ $[(A \text{ ind}_i \varphi) \wedge (A \text{ ind}_i \psi)] \Rightarrow A \text{ ind}_i(\varphi \wedge \psi)$
- ▶ $[(A \text{ ind}_i[A' \text{ holds}]) \wedge (A' \text{ ind}_i x)] \Rightarrow A \text{ ind}_i \varphi$
- ▶ $[(A \text{ ind}_i \varphi) \wedge (\varphi \text{ entails } \psi)] \Rightarrow A \text{ ind}_i \psi$

Some Properties

- ▶ $[(A \text{ holds}) \text{ entails } (A' \text{ holds})] \Rightarrow A \text{ ind}_i(A' \text{ holds})$
- ▶ $[(A \text{ ind}_i \varphi) \wedge (A \text{ ind}_i \psi)] \Rightarrow A \text{ ind}_i(\varphi \wedge \psi)$
- ▶ $[(A \text{ ind}_i[A' \text{ holds}]) \wedge (A' \text{ ind}_i x)] \Rightarrow A \text{ ind}_i \varphi$
- ▶ $[(A \text{ ind}_i \varphi) \wedge (\varphi \text{ entails } \psi)] \Rightarrow A \text{ ind}_i \psi$
- ▶ $[(A \text{ ind}_i R_j[A' \text{ holds}]) \wedge R_j(A' \text{ ind}_j \varphi)] \Rightarrow A \text{ ind}_i R_j(\varphi)$

Reflexive Common Indicator

Reflexive Common Indicator

- ▶ $A \text{ holds} \Rightarrow R_i(A \text{ holds})$

Reflexive Common Indicator

- ▶ $A \text{ holds} \Rightarrow R_i(A \text{ holds})$
- ▶ $A \text{ ind}_i R_j(A \text{ holds})$

Reflexive Common Indicator

- ▶ $A \text{ holds} \Rightarrow R_i(A \text{ holds})$
- ▶ $A \text{ ind}_i R_j(A \text{ holds})$
- ▶ $A \text{ ind}_i \varphi$

Reflexive Common Indicator

- ▶ $A \text{ holds} \Rightarrow R_i(A \text{ holds})$
- ▶ $A \text{ ind}_i R_j(A \text{ holds})$
- ▶ $A \text{ ind}_i \varphi$
- ▶ $(A \text{ ind}_i \psi) \Rightarrow R_i[A \text{ ind}_j \psi]$

Let $R^G(\varphi)$: $R_i\varphi, R_j\varphi, \dots, R_i(R_j\varphi), R_j(R_i(\varphi)), \dots$
iterated reason to believe φ .

Let $R^G(\varphi)$: $R_i\varphi, R_j\varphi, \dots, R_i(R_j\varphi), R_j(R_i(\varphi)), \dots$
iterated reason to believe φ .

Theorem. (Lewis) For all states of affairs A , for all propositions φ , and for all groups G : if A holds, and if A is a reflexive common indicator in G that φ , then $R^G(\varphi)$ is true.

Lewis and Aumann

Lewis common knowledge that φ *implies* the iterated definition of common knowledge (“Aumann common knowledge”)

Lewis and Aumann

Lewis common knowledge that φ *implies* the iterated definition of common knowledge (“Aumann common knowledge”), but the converse is not generally true....

Lewis and Aumann

Lewis common knowledge that φ *implies* the iterated definition of common knowledge (“Aumann common knowledge”), but the converse is not generally true....

Example. Suppose there is an agent $i \notin G$ that is *authoritative* for each member of G .

Lewis and Aumann

Lewis common knowledge that φ *implies* the iterated definition of common knowledge (“Aumann common knowledge”), but the converse is not generally true....

Example. Suppose there is an agent $i \notin G$ that is *authoritative* for each member of G . So, for $j \in G$, “ i states to j that φ is true” *indicates to j that φ .*

Lewis and Aumann

Lewis common knowledge that φ *implies* the iterated definition of common knowledge (“Aumann common knowledge”), but the converse is not generally true....

Example. Suppose there is an agent $i \notin G$ that is *authoritative* for each member of G . So, for $j \in G$, “ i states to j that φ is true” *indicates to j that φ* . Suppose that *separately and privately* to each member of G , i states that φ and $R^G(\varphi)$ are true.

Lewis and Aumann

Lewis common knowledge that φ *implies* the iterated definition of common knowledge (“Aumann common knowledge”), but the converse is not generally true....

Example. Suppose there is an agent $i \notin G$ that is *authoritative* for each member of G . So, for $j \in G$, “ i states to j that φ is true” *indicates to j that φ* . Suppose that *separately and privately* to each member of G , i states that φ and $R^G(\varphi)$ are true. Then, we have $R^i\varphi$ and $R_i(R^G(\varphi))$ for each $i \in G$.

Lewis and Aumann

Lewis common knowledge that φ *implies* the iterated definition of common knowledge (“Aumann common knowledge”), but the converse is not generally true....

Example. Suppose there is an agent $i \notin G$ that is *authoritative* for each member of G . So, for $j \in G$, “ i states to j that φ is true” *indicates to j that φ* . Suppose that *separately and privately* to each member of G , i states that φ and $R^G(\varphi)$ are true. Then, we have $R^i\varphi$ and $R_i(R^G(\varphi))$ for each $i \in G$. *But there is no common indicator that φ is true.*

Lewis and Aumann

Lewis common knowledge that φ *implies* the iterated definition of common knowledge (“Aumann common knowledge”), but the converse is not generally true....

Example. Suppose there is an agent $i \notin G$ that is *authoritative* for each member of G . So, for $j \in G$, “ i states to j that φ is true” *indicates to j that φ* . Suppose that *separately and privately* to each member of G , i states that φ and $R^G(\varphi)$ are true. Then, we have $R^i\varphi$ and $R_i(R^G(\varphi))$ for each $i \in G$. *But there is no common indicator that φ is true.* The agents $j \in G$ may have no reason to believe that everyone heard the statement from i or that all agents in G treat i as authoritative.

How does this help?

		<i>B</i>	
		<i>l</i>	<i>r</i>
<i>A</i>	<i>l</i>	10,10	0,0
	<i>r</i>	0,0	11,11

A: What should *we* do? **Team Reasoning**: why should this “mode of reasoning” be endorsed?

Reason to Believe

$R_i(\varphi)$: “agent i has reason to believe φ ” (deductive, inductive, norm of practical reason)

Reason to Believe

$R_i(\varphi)$: “agent i has reason to believe φ ” (deductive, inductive, norm of practical reason)

If i is the **subject of the proposition** φ_i , consider

$$R_i(\varphi_i) \text{ vs. } R_j(\varphi_i)$$

Reason to Believe

$R_i(\varphi)$: “agent i has reason to believe φ ” (deductive, inductive, norm of practical reason)

If i is the **subject of the proposition** φ_i , consider

$$R_i(\varphi_i) \text{ vs. } R_j(\varphi_i)$$

$$\text{inf}(R) : x, y \rightarrow z$$

$$\text{inf}(R) : x_1, \dots, x_n, \neg(x_1, \dots, x_n, \neg z) \rightarrow z$$

Common Reason to Believe

Awareness of Common Reason: for all $i \in G$ and all propositions φ ,

$$R^G(\varphi) \Rightarrow R_i[R^G(\varphi)]$$

Common Reason to Believe

Awareness of Common Reason: for all $i \in G$ and all propositions φ ,

$$R^G(\varphi) \Rightarrow R_i[R^G(\varphi)]$$

Authority of Common Reason: for all $i \in G$ and all propositions φ for which i is not the subject

$$\text{inf}(R_i) : R^G(\varphi) \rightarrow \varphi$$

Common Reason to Believe

Awareness of Common Reason: for all $i \in G$ and all propositions φ ,

$$R^G(\varphi) \Rightarrow R_i[R^G(\varphi)]$$

Authority of Common Reason: for all $i \in G$ and all propositions φ for which i is not the subject

$$\text{inf}(R_i) : R^G(\varphi) \rightarrow \varphi$$

Common Attribution of Common Reason: for all $i \in G$, for all propositions φ for which i is not the subject

$$\text{inf}(R^G) : \varphi \rightarrow R_i(\varphi)$$

Common Reason to Believe to Common Belief

Theorem The three previous properties can generate any hierarchy of belief (i has reason to believe that j has reason to believe that... that φ) for any φ with $R^G(\varphi)$.

Team Maximising

$inf(R_i) : R^N[opt(v, N, s^N)],$
 $R^N[\text{each } i \in N \text{ endorses team maximising with respect to } N \text{ and } v],$
 $R^N[\text{each member of } N \text{ acts on reasons }] \rightarrow ought(i, s_i)$

Team Maximising

$inf(R_i) : R^N[opt(v, N, s^N)],$

$R^N[\text{each } i \in N \text{ endorses team maximising with respect to } N \text{ and } v],$

$R^N[\text{each member of } N \text{ acts on reasons }] \rightarrow \textit{ought}(i, s_i)$

$R_i[\textit{ought}(i, s_i)]: i \text{ has reason to choose } s_i$

Team Maximising

$inf(R_i) : R^N[opt(v, N, s^N)],$
 $R^N[\text{each } i \in N \text{ endorses team maximising with respect to } N \text{ and } v],$
 $R^N[\text{each member of } N \text{ acts on reasons }] \rightarrow ought(i, s_i)$

$i \text{ acts on reasons if for all } s_i, R[ought(i, s_i)] \Rightarrow choice(i, s_i)$

Team Maximising

$inf(R_i) : R^N[opt(v, N, s^N)],$
 $R^N[\text{each } i \in N \text{ endorses team maximising with respect to } N \text{ and } v],$
 $R^N[\text{each member of } N \text{ acts on reasons }] \rightarrow ought(i, s_i)$

$opt(v, N, s^N)$: s^N is maximal for the group N w.r.t. v

Team Maximising

$inf(R_i) : R^N[opt(v, N, s^N)],$
 $R^N[\text{each } i \in N \text{ endorses team maximising with respect to } N \text{ and } v],$
 $R^N[\text{each member of } N \text{ acts on reasons}] \rightarrow ought(i, s_i)$

Next

Team modes of reasoning, group identification, frames and team preferences

Levels of Knowledge

Fix a set of agents $\mathcal{A} = \{1, \dots, n\}$.

$\Sigma_K = \{K_1, \dots, K_n\}$ and $\Sigma_C = \{C_U\}_{U \subseteq \mathcal{A}}$

Level of Knowledge: $Lev_{\mathcal{M}}(p, s) = \{x \in \Sigma^* \mid \mathcal{M}, s \models xp\}$
(where $\Sigma = \Sigma_K$ or $\Sigma = \Sigma_C$).

[If Σ is a finite set, then Σ^* is the set of finite strings over Σ]
[Recall the definition of truth in a Kripke structure]

Levels of Knowledge

Fix a set of agents $\mathcal{A} = \{1, \dots, n\}$.

$\Sigma_K = \{K_1, \dots, K_n\}$ and $\Sigma_C = \{C_U\}_{U \subseteq \mathcal{A}}$

Level of Knowledge: $Lev_{\mathcal{M}}(p, s) = \{x \in \Sigma^* \mid \mathcal{M}, s \models xp\}$
(where $\Sigma = \Sigma_K$ or $\Sigma = \Sigma_C$).

[If Σ is a finite set, then Σ^* is the set of finite strings over Σ]
[Recall the definition of truth in a Kripke structure]

R. Parikh and P. Krasucki. *Levels of knowledge in distributed computing*. Sadhana-Proceedings of the Indian Academy of Science 17 (1992).

R. Parikh. *Levels of knowledge, games and group action*. Research in Economics 57, pp. 267 - 281 (2003).

Levels of Knowledge

A level of knowledge is simply a set of finite strings over Σ_C .
Why isn't it obvious that there are *uncountably* many levels of knowledge?

Levels of Knowledge

A level of knowledge is simply a set of finite strings over Σ_C .
Why isn't it obvious that there are *uncountably* many levels of knowledge?

Consider the sets:

- ▶ $L_1 = \{K_1, K_2\}$ and $L_2 = \{K_1, K_2, K_1K_2\}$

- ▶ $L_1 = \{K_1, K_3, K_1K_2K_3\}$ and $L_2 = \{K_1, K_2, K_3, K_1K_2K_3\}$

Levels of Knowledge

A level of knowledge is simply a set of finite strings over Σ_C .
Why isn't it obvious that there are *uncountably* many levels of knowledge?

Consider the sets:

- ▶ $L_1 = \{K_1, K_2\}$ and $L_2 = \{K_1, K_2, K_1K_2\}$
(*different level of knowledge*)
- ▶ $L_1 = \{K_1, K_3, K_1K_2K_3\}$ and $L_2 = \{K_1, K_2, K_3, K_1K_2K_3\}$
(*same level of knowledge*)

Levels of Knowledge: Preliminaries

Given any two strings $x, y \in \Sigma^*$, we say x is **embeddable** in y , written $x \leq y$, if all symbols of x occur in y in the same order but not necessarily consecutively.

Levels of Knowledge: Preliminaries

Given any two strings $x, y \in \Sigma^*$, we say x is **embeddable** in y , written $x \leq y$, if all symbols of x occur in y in the same order but not necessarily consecutively. Formally, we can define \leq as follows:

1. $x \leq x$ and $\epsilon \leq x$ for all $x \in \Sigma^*$
2. $x \leq y$ if there exists x', x'', y', y'' , ($y, y'' \neq \epsilon$) such that $x = x'x''$, $y = y'y''$ and $x' \leq y'$, $x'' \leq y''$.

\leq is the smallest relation satisfying (1) and (2).

Levels of Knowledge: Preliminaries

Given any two strings $x, y \in \Sigma^*$, we say x is **embeddable** in y , written $x \leq y$, if all symbols of x occur in y in the same order but not necessarily consecutively. Formally, we can define \leq as follows:

1. $x \leq x$ and $\epsilon \leq x$ for all $x \in \Sigma^*$
2. $x \leq y$ if there exists x', x'', y', y'' , ($y, y'' \neq \epsilon$) such that $x = x'x''$, $y = y'y''$ and $x' \leq y'$, $x'' \leq y''$.

\leq is the smallest relation satisfying (1) and (2).

Example:

$aba \leq aaba$

$aba \leq abca$

$aba \not\leq aabb$

Levels of Knowledge: Preliminaries

(X, \preceq) is

Levels of Knowledge: Preliminaries

(X, \preceq) is

a **partial order** if \preceq is reflexive, transitive and antisymmetric.

well founded if every infinite subset of X has a $(\preceq -)$ minimal element.

a **linear order** if it is a partial order and all elements of X are comparable.

Levels of Knowledge: Preliminaries

(X, \preceq) is

a **partial order** if \preceq is reflexive, transitive and antisymmetric.

well founded if every infinite subset of X has a $(\preceq -)$ minimal element.

a **linear order** if it is a partial order and all elements of X are comparable.

a **well-partial order** (WPO) if (X, \preceq) is a partial order and every linear order that extends (X, \preceq) (i.e., a linear order (X, \preceq') with $\preceq \subseteq \preceq'$) is well-founded.

Levels of Knowledge: Preliminaries

(X, \preceq) is

a **partial order** if \preceq is reflexive, transitive and antisymmetric.

well founded if every infinite subset of X has a $(\preceq -)$ minimal element.

a **linear order** if it is a partial order and all elements of X are comparable.

a **well-partial order** (WPO) if (X, \preceq) is a partial order and every linear order that extends (X, \preceq) (i.e., a linear order (X, \preceq') with $\preceq \subseteq \preceq'$) is well-founded.

A set $\{a_1, a_2, \dots\}$ of incomparable elements is a well-founded partial order but not a WPO.

Well-Partial Orders

Fact. (X, \preceq) is a WPO iff \preceq is well-founded and every subset of mutually incomparable elements is finite

Well-Partial Orders

Fact. (X, \preceq) is a WPO iff \preceq is well-founded and every subset of mutually incomparable elements is finite

Theorem (Higman). If Σ is finite, then (Σ^*, \leq) is a WPO

G. Higman. *Ordering by divisibility in abstract algebras*. Proc. London Math. Soc. 3 (1952).

D. de Jongh and R. Parikh. *Well-Partial Orderings and Hierarchies*. Proc. of the Koninklijke Nederlandse Akademie van Wetenschappen 80 (1977).

WPO and Downward Closed Sets

Given (X, \preceq) a set $A \subseteq X$ is **downward closed** iff $x \in A$ implies for all $y \preceq x$, $y \in A$.

Theorem. (Parikh & Krasucki) If Σ is finite, then there are only countably many \preceq -downward closed subsets of Σ^* and all of them are *regular*.

Levels of Knowledge

Theorem. Consider the alphabet $\Sigma_C = \{C_U\}_{U \subseteq \mathcal{A}}$. For all strings $x, y \in \Sigma_C^*$, if $x \preceq y$ then for all pointed models \mathcal{M}, s , if $\mathcal{M}, s \models yP$ then $\mathcal{M}, s \models xP$.

Levels of Knowledge

Theorem. Consider the alphabet $\Sigma_C = \{C_U\}_{U \subseteq \mathcal{A}}$. For all strings $x, y \in \Sigma_C^*$, if $x \preceq y$ then for all pointed models \mathcal{M}, s , if $\mathcal{M}, s \models yP$ then $\mathcal{M}, s \models xP$.

Incorrect as stated, but easily fixed: every extension of a WPO is a WPO

1. $K_1 K_1 \not\preceq K_1$
2. We should have $C_U \preceq C_V$ if $U \subseteq V$.

Levels of Knowledge

Theorem. Consider the alphabet $\Sigma_C = \{C_U\}_{U \subseteq \mathcal{A}}$. For all strings $x, y \in \Sigma_C^*$, if $x \preceq y$ then for all pointed models \mathcal{M}, s , if $\mathcal{M}, s \models yP$ then $\mathcal{M}, s \models xP$.

Incorrect as stated, but easily fixed: every extension of a WPO is a WPO

1. $K_1 K_1 \not\preceq K_1$
2. We should have $C_U \preceq C_V$ if $U \subseteq V$.

Corollary 1. Every level of knowledge is a downward closed set.

Levels of Knowledge

Theorem. Consider the alphabet $\Sigma_C = \{C_U\}_{U \subseteq \mathcal{A}}$. For all strings $x, y \in \Sigma_C^*$, if $x \preceq y$ then for all pointed models \mathcal{M}, s , if $\mathcal{M}, s \models yP$ then $\mathcal{M}, s \models xP$.

Incorrect as stated, but easily fixed: every extension of a WPO is a WPO

1. $K_1 K_1 \not\preceq K_1$
2. We should have $C_U \preceq C_V$ if $U \subseteq V$.

Corollary 1. Every level of knowledge is a downward closed set.

Corollary 2. There are only countably many levels of knowledge.

Realizing Levels of Knowledge

Theorem. (R. Parikh and EP) Suppose that L is a downward closed subset of Σ_K^* , then there is a finite Kripke model \mathcal{M} and state s such that $\mathcal{M}, s \models xP$ iff $x \in L$. (i.e., $L = Lev_{\mathcal{M}}(p, s)$).

▶ Back

States of Knowledge

S. Hart, A. Heifetz and D. Samet. *"Knowing Whether," "Knowing That," and The Cardinality of State Spaces*. Journal of Economic Theory 70 (1996).

States of Knowledge

Let W be a set of states and fix an event $X \subseteq W$.

Consider a sequence of finite boolean algebras $\mathcal{B}_0, \mathcal{B}_1, \mathcal{B}_2, \dots$ defined as follows:

$$\mathcal{B}_0 = \{\emptyset, X, \neg X, \Omega\}$$

$$\mathcal{B}_n = \mathcal{B}_{n-1} \cup \{K_i E \mid E \in \mathcal{B}_{n-1}, i \in \mathcal{A}\}$$

The events $\mathcal{B} = \cup_{i=1,2,\dots} \mathcal{B}_i$ are said to be **generated by** X .

States of Knowledge

Definition. Two states w, w' are **separated** by X if there exists an event E which is generated by X such that $w \in E$ and $w' \in \neg E$.

Question: How many states can be in an information structure (W, Π_1, Π_2) such that an event X separates any two of them?

States of Knowledge

Consider a K -list (E_1, E_2, E_3, \dots) of events generated by X .

We can of course, write down infinitely many infinite K -lists (**uncountably many!**).

States of Knowledge

Consider a K -list (E_1, E_2, E_3, \dots) of events generated by X .

We can of course, write down infinitely many infinite K -lists (**uncountably many!**).

Again, are they all consistent?

States of Knowledge

Consider a K -list (E_1, E_2, E_3, \dots) of events generated by X .

We can of course, write down infinitely many infinite K -lists (**uncountably many!**).

Again, are they all consistent?

Consider $(X, K_1X, \neg K_2K_1X, \neg K_1\neg K_2K_1X, K_2\neg K_1\neg K_2K_1X)$

States of Knowledge

Consider a K -list (E_1, E_2, E_3, \dots) of events generated by X .

We can of course, write down infinitely many infinite K -lists (**uncountably many!**).

Again, are they all consistent?

$(X, K_1X, \neg K_2K_1X, \neg K_1\neg K_2K_1X, K_2\neg K_1\neg K_2K_1X)$ is inconsistent.

Knowing Whether

Let $J_i E := K_i E \vee K_i \neg E$.

Lemma. Every J -list is consistent.

Theorem. (Hart, Heifetz and Samet) There are uncountably many states of knowledge.

S. hart, A. Heifetz and D. Samet. *"Knowing Whether," "Knowing That," and The Cardinality of State Spaces.* Journal of Economic Theory 70 (1996).

What about beliefs?

In Aumann/Kripke structures belief operators are just like knowledge operators except we replace the truth axiom/property ($K\varphi \rightarrow \varphi$) with a consistency property ($\neg B\perp$).

What about beliefs?

In Aumann/Kripke structures belief operators are just like knowledge operators except we replace the truth axiom/property ($K\varphi \rightarrow \varphi$) with a consistency property ($\neg B\perp$).

Theorem. (R. Parikh and EP) There are uncountably many levels of belief.

▶ Back

The Issue

	<i>L</i>	<i>R</i>
<i>U</i>	1,1	0,1
<i>D</i>	0,2	1,0

The Issue

	L	R
U	1,1	0,1
D	0,2	1,0

Suppose rationality incorporates *admissibility* (or *cautiousness*).

The Issue

	L	R
U	1,1	0,1
D	0,2	1,0

Suppose rationality incorporates *admissibility* (or *cautiousness*).

1. Both Row and Column should use a *full-support* probability measure
2. But if Row thinks that Column is **rational** then should she not assign probability 1 to L ?

The Issue

	<i>L</i>	<i>R</i>
<i>U</i>	1,1	0,1
<i>D</i>	0,2	1,0

Suppose rationality incorporates *admissibility* (or *cautiousness*).

1. Both Row and Column should use a *full-support* probability measure
2. But if Row thinks that Column is **rational** then should she not assign probability 1 to *L*?

The condition that the players incorporate admissibility into their rationality calculations seems to conflict with the condition that the players think the other players are rational

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

L. Samuelson. *Dominated Strategies and Common Knowledge*. Games and Economic Behavior (1992).

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

	Y_1	Y_2	Y_3
X_1	2,4	5,4	-1,0
X_2	3,4	2,4	-2,0
X_3	1,2	0,0	2,2
X_4	0,2	2,0	0,4

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

	Y_1	Y_2	Y_3
X_1	2,4	5,4	-1,0
X_2	3,4	2,4	-2,0
X_3	1,2	0,0	2,2
X_4	0,2	2,0	0,4

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

	Y_1	Y_3
X_1	2,4	-1,0
X_2	3,4	-2,0
X_3	1,2	2,2
X_4	0,2	0,4

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

	Y_1	Y_3
X_1	2,4	-1,0
X_2	3,4	-2,0
X_3	1,2	2,2
X_4	0,2	0,4

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

	Y_1	Y_3
X_1	2,4	-1,0
X_2	3,4	-2,0
X_3	1,2	2,2

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

	Y_1	Y_3
X_1	2,4	-1,0
X_2	3,4	-2,0
X_3	1,2	2,2

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

	Y_1
X_1	2,4
X_2	3,4
X_3	1,2

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

	Y_1
X_2	3,4

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

	Y_1	Y_2	Y_3
X_1	2,4	5,4	-1,0
X_2	3,4	2,4	-2,0
X_3	1,2	0,0	2,2
X_4	0,2	2,0	0,4

$\{X_2, Y_1\}$ is the unique IA solution, but common knowledge of admissibility yields a unique *consistent pair*:
 $\{\Delta(X_1, X_2), \Delta(Y_1, Y_2)\}$.

Common Knowledge of Admissibility

Theorem Iterated admissibility is not equivalent to common knowledge of admissibility.

	Y_1	Y_2	Y_3
X_1	2,4	5,4	-1,0
X_2	3,4	2,4	-2,0
X_3	1,2	0,0	2,2
X_4	0,2	2,0	0,4

$\{X_2, Y_1\}$ is the unique IA solution, but common knowledge of admissibility yields a unique *consistent pair*:
 $\{\Delta(X_1, X_2), \Delta(Y_1, Y_2)\}$.

Other Results

1. We have seen that IA and common knowledge of admissibility diverge.

L. Samuelson. *Dominated Strategies and Common Knowledge*. Games and Economic Behavior (1992).

Other Results

1. We have seen that IA and common knowledge of admissibility diverge.
2. There exist games in which assuming that admissibility is common knowledge does not provide players with sufficient information to determine which strategies should be eliminated on admissibility grounds.

L. Samuelson. *Dominated Strategies and Common Knowledge*. Games and Economic Behavior (1992).

Other Results

1. We have seen that IA and common knowledge of admissibility diverge.
2. There exist games in which assuming that admissibility is common knowledge does not provide players with sufficient information to determine which strategies should be eliminated on admissibility grounds.
3. There exists games in which assuming that admissibility is common knowledge yields a contradiction

L. Samuelson. *Dominated Strategies and Common Knowledge*. Games and Economic Behavior (1992).

Both Including and Excluding a Strategy

One solution is to assume that players consider some strategies *infinitely more likely than other strategies*.

Both Including and Excluding a Strategy

One solution is to assume that players consider some strategies *infinitely more likely than other strategies*.

Lexicographic Probability System: a sequence of probability distributions each infinitely more likely than the next.

Both Including and Excluding a Strategy

One solution is to assume that players consider some strategies *infinitely more likely than other strategies*.

Lexicographic Probability System: a sequence of probability distributions each infinitely more likely than the next.

	1	[1]
	<i>L</i>	<i>R</i>
<i>U</i>	1,1	0,1
<i>D</i>	0,2	1,0

A. Brandenburger, A. Friedenberg, H. J. Keisler. *Admissibility in Games*. *Econometrica* (2008).

▶ Back