

Claude's Cycles

Don Knuth, Stanford Computer Science Department
(28 February 2026; revised 02 March 2026)

Shock! Shock! I learned yesterday that an open problem I'd been working on for several weeks had just been solved by Claude Opus 4.6—Anthropic's hybrid reasoning model that had been released three weeks earlier! It seems that I'll have to revise my opinions about “generative AI” one of these days. What a joy it is to learn not only that my conjecture has a nice solution but also to celebrate this dramatic advance in automatic deduction and creative problem solving. I'll try to tell the story briefly in this note.

Here's the problem, which came up while I was writing about directed Hamiltonian cycles for a future volume of *The Art of Computer Programming*:

Consider the digraph with m^3 vertices ijk for $0 \leq i, j, k < m$, and three arcs from each vertex, namely to i^+jk , ij^+k , and ijk^+ , where $i^+ = (i+1) \bmod m$. Try to find a general decomposition of the arcs into three directed m^3 -cycles, for all $m > 2$.

I had solved the problem for $m = 3$, and asked for a generalization as part of the answer to an exercise in [3]. My friend Filip Stappers rose to the challenge, and empirically discovered solutions for $4 \leq m \leq 16$; therefore it became highly likely that the desired decompositions do exist, except when $m \leq 2$.

Indeed, it was Filip who had the gumption to pose this question to Claude, using exactly the wording above. He also gave guidance/coaching, instructing Claude to summarize its ongoing progress:

```
** After EVERY exploreXX.py run, IMMEDIATELY update this file [plan.md]
before doing anything else. ** No exceptions. Do not start the next exploration
until the previous one is documented here.
```

And Claude's plan of attack was quite admirable. First it reformulated the problem: “Need sigma: $Z_m^3 \rightarrow S_3$, assigning a permutation of $\{0, 1, 2\}$ at each vertex; cycle c at vertex v goes in direction $\text{sigma}(v)[c]$. Each resulting functional digraph must be a single Hamiltonian cycle.” Then it tried for a simple answer: “If sigma is cyclic, i.e. $f_c(v) = (c+g(v)) \bmod 3$ for some $g: Z_m^3 \rightarrow Z_3$, then we only need one function g . Tested linear $g = (a \cdot i + b \cdot j + c \cdot k) \bmod 3$ and quadratic variants. *No linear or simple quadratic g works.*”

Claude's second exploration was what it called a DFS search attempt (depth-first search): “Attempted brute-force DFS over sigma assignments for $m = 3$ (6^{27} search space). Too slow without good pruning.”

Thirdly, it tried what it called 2D serpentine analysis: “*Key 2D result:* The pattern $[0^{m-1}1]^m$ gives a Hamiltonian cycle in $\text{Cay}(Z_m^2, \{e_0, e_1\})$ for $m = 3$. Its complement also works, giving a valid 2D decomposition.” This is really impressive! First Claude recognized the digraph as a so-called Cayley digraph with two generators, traditionally called e_0 and e_1 ; a mathematician will understand that the arcs act as a group on m^2 vertices, where e_0 increments the first component while e_1 increments the second. Then it formulated the cycle symbolically for general m , although for some reason it still said that it was only looking at $m = 3$. And it decided to call this a “serpentine pattern”!

Exploration number 4 constructed the “3D serpentine pattern” $[0^{m-1}1]^{m-1}[0^{m-1}2]^m$. (That's a typographic error for $[[0^{m-1}1]^{m-1}0^{m-1}2]^m$. It's a classical sequence called the “modular m -ary Gray code” on page 299 of [2].) After deleting this cycle from the graph, Claude looked at the remaining 2-regular digraph and found that it has “rigid structure making decomposition difficult.”

Nevertheless, exploration number 5 analyzed that residual digraph, finding some cycles. Unfortunately they didn't work out.

Back to the drawing board. Several other approaches were tried, for example looking at the choices to be made at vertices that lie on the hyperplane where $i+j+k$ is congruent to -1 . Nothing promising showed up until exploration number 15, which introduced what Claude called a *fiber decomposition*: “The quotient map $?(i, j, k) = i+j+k \bmod m$ maps all arcs from fiber F_s to F_{s+1} . The digraph is *layered*. In fiber coordinates (i, j) with $k = (s-i-j) \bmod m$.”

Using those fiber coordinates, the problem could now be reformulated as choosing, for each (i, j, s) , a permutation of the three operators that shift (i, j) by either $(1, 0)$ or $(0, 1)$ or $(0, 0)$. “Tried sigma depending

only on fiber index s (rotation by s): gives the $(i+j+k) \bmod 3$ pattern which doesn't work. Random local search found solutions for $m = 3$ but no general pattern emerged."

Exploration 18 "implemented the full fiber framework as executable code. *Exhaustive backtracking* for $m = 3$ found a valid decomposition in 0.1s (10932 nodes)." In this solution the permutation sigma was the same for all (i, j) in fiber $s = 0$. A similarly "uniform" fiber was found for $m = 4$ by simulated annealing, as part of exploration 20.

In exploration 21, Claude realized that the 2D serpentine function " $Q(i, j) = (i+1, j)$ if $i+j \neq m-1 \bmod m$, else $(i, j+1)$ " had nice properties. It looked promising when a uniform choice is made at fiber 0 and a version of the serpentine at all other fibers, if and only if m is odd.

That triggered a bunch of experiments with simulated annealing "at scale"; but all was for naught. The conclusion after exploration 25 was "SA can find solutions but cannot give a general construction. Need pure math."

So now Claude put on a new thinking cap. At one point it said to itself, "Maybe the right framing is: don't think in fibers, think directly about what makes a Hamiltonian cycle."

There was a near miss at exploration number 27. "Take the 3D serpentine for cycle 0, apply cyclic coordinate rotation to get cycles 1 and 2: $d_1(i, j, k) = (d_0(j, k, i) + 1) \bmod 3$; $d_2(i, j, k) = (d_0(k, i, j) + 2) \bmod 3$; *all three are individually Hamiltonian* for every $m = 3..9$ tested! Only $3(m-1)$ out of m^3 vertices have conflicts. *all* lie on the hyperplane $i+j+k = m-1 \bmod m$."

Unfortunately, those conflicts couldn't be resolved. Exploration 29 proved that many plausible scenarios were in fact impossible. "This kills the 'single-hyperplane + rotation' approach entirely. . . . We must allow the direction function to use *different* values across a rotation orbit."

But, aha. Exploration 30 went back to the solution found by SA in exploration 20 and noticed that the choice at each fiber depends on only a single coordinate: $s = 0$ only on j ; $s = 1$ and $s = 2$ only on i . This led to a concrete construction (exploration 31), in the form of a Python program, which produced valid results for $m = 3, 5, 7, 9$, and 11—hooray! "All three cycles are Hamiltonian, all arcs are used, perfect decomposition!"

Here is a simplification of that Python program, converted into C form:

```

int c, i, j, k, m, s, t;
char *d;
for (c = 0; c < 3; c++) {
    for (t = i = j = k = 0; ; t++) {
        printf ("%x%x%x ", i, j, k);
        if (t == m*m*m) break;
        s = (i+j+k) % m;
        if (s == 0) d = (j == m-1? "012" : "210");
        else if (s == m-1) d = (i > 0? "120" : "210");
        else d = (i == m-1? "201" : "102");
        switch (d[c]) {
        case '0': i = (i+1) % m; break;
        case '1': j = (j+1) % m; break;
        case '2': k = (k+1) % m; break;
        }
    }
    printf ("\n");
}

```

Filip Stappers tested Claude's Python program for all odd m between 3 and 101, finding perfect decompositions each time. Thus he concluded, quite reasonably, that the problem was indeed solved for odd values of m . And he quickly sent me the shocking news.

Of course, a rigorous proof was still needed. And the construction of such a proof turns out to be quite interesting. Let's look, for example, at the first cycle that is printed; we must prove that it has length m^3 .

The rule for that cycle is nontrivial, yet fairly simple: Let $s = (i+j+k) \bmod m$.

When $s = 0$, bump i if $j = m-1$, otherwise bump k .
 When $0 < s < m-1$, bump k if $i = m-1$, otherwise bump j .
 When $s = m-1$, bump j if $i > 0$, otherwise bump k .

(“Bump” means “increase by 1, mod m .”) Hence in the special case $m = 3$ that cycle is

022 \rightarrow 002 \rightarrow 000 \rightarrow 001 \rightarrow 011 \rightarrow 012 \rightarrow 010 \rightarrow 020 \rightarrow 021 \rightarrow
 121 \rightarrow 101 \rightarrow 111 \rightarrow 112 \rightarrow 122 \rightarrow 102 \rightarrow 100 \rightarrow 110 \rightarrow 120 \rightarrow
 220 \rightarrow 221 \rightarrow 201 \rightarrow 202 \rightarrow 200 \rightarrow 210 \rightarrow 211 \rightarrow 212 \rightarrow 222 \rightarrow 022.

And in the special case $m = 5$ it's

042 \rightarrow 002 \rightarrow 012 \rightarrow 022 \rightarrow 023 \rightarrow 024 \rightarrow 034 \rightarrow 044 \rightarrow 004 \rightarrow 000 \rightarrow 001 \rightarrow 011 \rightarrow 021 \rightarrow
 031 \rightarrow 032 \rightarrow 033 \rightarrow 043 \rightarrow 003 \rightarrow 013 \rightarrow 014 \rightarrow 010 \rightarrow 020 \rightarrow 030 \rightarrow 040 \rightarrow 041 \rightarrow
 141 \rightarrow 101 \rightarrow 111 \rightarrow 121 \rightarrow 131 \rightarrow 132 \rightarrow 142 \rightarrow 102 \rightarrow 112 \rightarrow 122 \rightarrow 123 \rightarrow 133 \rightarrow 143 \rightarrow
 103 \rightarrow 113 \rightarrow 114 \rightarrow 124 \rightarrow 134 \rightarrow 144 \rightarrow 104 \rightarrow 100 \rightarrow 110 \rightarrow 120 \rightarrow 130 \rightarrow 140 \rightarrow
 240 \rightarrow 200 \rightarrow 210 \rightarrow 220 \rightarrow 230 \rightarrow 231 \rightarrow 241 \rightarrow 201 \rightarrow 211 \rightarrow 221 \rightarrow 222 \rightarrow 232 \rightarrow 242 \rightarrow
 202 \rightarrow 212 \rightarrow 213 \rightarrow 223 \rightarrow 233 \rightarrow 243 \rightarrow 203 \rightarrow 204 \rightarrow 214 \rightarrow 224 \rightarrow 234 \rightarrow 244 \rightarrow
 344 \rightarrow 304 \rightarrow 314 \rightarrow 324 \rightarrow 334 \rightarrow 330 \rightarrow 340 \rightarrow 300 \rightarrow 310 \rightarrow 320 \rightarrow 321 \rightarrow 331 \rightarrow 341 \rightarrow
 301 \rightarrow 311 \rightarrow 312 \rightarrow 322 \rightarrow 332 \rightarrow 342 \rightarrow 302 \rightarrow 303 \rightarrow 313 \rightarrow 323 \rightarrow 333 \rightarrow 343 \rightarrow
 443 \rightarrow 444 \rightarrow 440 \rightarrow 441 \rightarrow 401 \rightarrow 402 \rightarrow 403 \rightarrow 404 \rightarrow 400 \rightarrow 410 \rightarrow 411 \rightarrow 412 \rightarrow 413 \rightarrow
 414 \rightarrow 424 \rightarrow 420 \rightarrow 421 \rightarrow 422 \rightarrow 423 \rightarrow 433 \rightarrow 434 \rightarrow 430 \rightarrow 431 \rightarrow 432 \rightarrow 442 \rightarrow 042.

Triples for the same value of s are spaced exactly m steps apart. To prove that all m^3 triples occur, we need to prove that the m^2 triples for any given value of s are all present.

Notice that the first coordinate, i , changes only when $s = 0$ and $j = m-1$. Thus the m^2 triples with any given value of the first coordinate i all occur consecutively. (Our example cycles indicate this by starting at the vertex where i has just changed to 0, instead of starting at vertex 000.)

Notice that the cycle elements with $i = 0$ must start in general with the vertex $0(m-1)2$, because the previous vertex must have had $i = j = m-1$ and $s = 0$. And $0(m-1)2$, which has $s = 1$, is followed in general by $002, 012, \dots, 0(m-3)2, 0(m-3)3$, taking us back to $s = 0$.

In general $0(m-k)k$ is immediately followed by $0(m-k)(k+1)$ when $1 < k < m$; and then j increases until we reach $0(m-k-2)(k+1)$, whose successor is $0(m-k-2)(k+2)$. Thus k is increased by 2, modulo m , each time we hit a vertex with $s = 0$. Since m is odd, we'll eventually get to $0(m-1)1$ —at which time we will have seen all m^2 vertices of the form $0jk$. And the successor of $0(m-1)1$ is $1(m-1)1$.

So far so good! In general, the cycle elements whose first component i satisfies $0 < i < m-1$ will start with $i(m-1)(2-i)$, where $2-i$ is of course evaluated mod m . If all goes well those elements should include all m^2 vertices that begin with i , ending with $i(m-i)(1-i)$ —whose successor is $(i+1)(m-1)(1-i)$. And all *does* go well: We repeatedly advance the second component except when $s = 0$; hence the vertices that we see when $s = 0$ are $i(m-2)(2-i), i(m-3)(3-i), \dots, i0(m-i), i(m-1)(1-i)$.

Finally we reach $(m-1)(m-1)3$, the first vertex for which $i = m-1$. The local rules change again: From now on we'll repeatedly advance the *third* component, except when $s = m-1$. The vertices we see when $s = 0$ are $(m-1)01, (m-1)10, \dots, (m-1)(m-1)2$. QED.

We've now proved that the first cycle (the cycle for $c = 0$ in the C program) is Hamiltonian. Similar proofs can be carried out for the other two cycles. (See the Appendix.)

For fun, let's consider a larger class of cycles for which such proofs exist. Indeed, it turns out that there are hundreds of way to solve the stated decomposition problem for odd m ; Claude Opus 4.6 happened to discover just one of them, by deducing where to look.

We shall say that a Hamiltonian cycle C for $m = 3$ is *generalizable* if the following construction yields a Hamiltonian cycle for all odd $m \geq 3$: “Given an arbitrary vertex IJK for $0 \leq I, J, K < m$, let $i = I'$, $j = J'$, $s = S'$, and $k = (s - i - j) \bmod 3$, where $S = (I + J + K) \bmod m$, $0' = 0$, $(m - 1)' = 2$, and $x' = 1$ for $0 < x < m - 1$. Obtain the successor of IJK by bumping the coordinate that cycle C bumps when forming the successor of ijk .” For example, if $m = 5$ and $IJK = 301$, we have $i = 1$, $j = 0$, $S = 4$, $s = 2$, $k = 1$. So the successor of 301 will be either 401 or 311 or 302, depending on whether C contains the arc $101 \rightarrow 201$ or $101 \rightarrow 111$ or $101 \rightarrow 102$.

Claude’s cycle in the example above for $m = 3$ is generalizable; indeed, we’ve seen its generalization for $m = 5$. But it’s easy to find cycles that are not generalizable. For instance, if C is the cycle

000 \rightarrow 001 \rightarrow 002 \rightarrow 012 \rightarrow 010 \rightarrow 011 \rightarrow 021 \rightarrow 022 \rightarrow 020 \rightarrow 120 \rightarrow 100 \rightarrow 101 \rightarrow 111 \rightarrow 121 \rightarrow
 122 \rightarrow 102 \rightarrow 112 \rightarrow 110 \rightarrow 210 \rightarrow 211 \rightarrow 212 \rightarrow 222 \rightarrow 220 \rightarrow 221 \rightarrow 201 \rightarrow 202 \rightarrow 200 \rightarrow 000

(which happens to be the lexicographically smallest of all Hamiltonian cycles for the case $m = 3$), we get the following “generalization” for $m = 5$:

000 \rightarrow 001 \rightarrow 002 \rightarrow 003 \rightarrow 004 \rightarrow 014 \rightarrow 010 \rightarrow 011 \rightarrow 012 \rightarrow 013 \rightarrow 023 \rightarrow 024 \rightarrow 020 \rightarrow
 021 \rightarrow 022 \rightarrow 032 \rightarrow 033 \rightarrow 034 \rightarrow 030 \rightarrow 031 \rightarrow 041 \rightarrow 042 \rightarrow 043 \rightarrow 044 \rightarrow 040 \rightarrow 140 \rightarrow
 100 \rightarrow 101 \rightarrow 102 \rightarrow 103 \rightarrow 113 \rightarrow 123 \rightarrow 124 \rightarrow 120 \rightarrow 121 \rightarrow 221 \rightarrow 231 \rightarrow 232 \rightarrow 233 \rightarrow
 234 \rightarrow 334 \rightarrow 344 \rightarrow 340 \rightarrow 341 \rightarrow 342 \rightarrow 302 \rightarrow 312 \rightarrow 313 \rightarrow 314 \rightarrow 310 \rightarrow 410 \rightarrow 411 \rightarrow
 412 \rightarrow 413 \rightarrow 414 \rightarrow 424 \rightarrow 420 \rightarrow 421 \rightarrow 422 \rightarrow 423 \rightarrow 433 \rightarrow 434 \rightarrow 430 \rightarrow 431 \rightarrow 432 \rightarrow
 442 \rightarrow 443 \rightarrow 444 \rightarrow 440 \rightarrow 441 \rightarrow 401 \rightarrow 402 \rightarrow 403 \rightarrow 404 \rightarrow 400 \rightarrow 000.

Oops — this cycle has length 75, not 125.

It turns out that there are exactly 11502 Hamiltonian cycles for $m = 3$, of which exactly 1012 generalize to Hamiltonian cycles for $m = 5$. Furthermore, exactly 996 of them generalize to Hamiltonian cycles for both $m = 5$ and $m = 7$. And those 996 are in fact generalizable to all odd $m > 1$.

Now here’s the point: Let’s say that a decomposition is “Claude-like” if it can be generated by a C program like the one above, in which the permutations d of $\{0, 1, 2\}$ depend only on whether i , j , and s are 0 or $m - 1$ or not. No special cases other than 0 and $m - 1$ are allowed to affect the choice of d .

Theorem. *A Claude-like decomposition is valid for all odd $m > 1$ if and only if each of the three sequences that it defines for $m = 3$ is generalizable.*

Proof. If those three sequences aren’t Hamiltonian, or if they don’t generalize to Hamiltonian cycles for all odd $m > 1$, they certainly don’t define a valid decomposition. Conversely, if they are Hamiltonian cycles that generalize to Hamiltonian cycles for all odd $m > 1$, they certainly are valid: Every vertex ijk appears in each of the three cycles, and its three outgoing arcs are partitioned properly because d is a permutation. ■

By setting up an exact cover problem, using the 11502 Hamiltonian cycles for $m = 3$, we can deduce that there are exactly 4554 solutions to the $3 \times 3 \times 3$ decomposition problem. And in a similar fashion, if we study all ways to cover every arc by using just three of the 996 cycles that are generalizable, we find that exactly 760 of those 4554 solutions involve only generalizable cycles — about one in every 6. Consequently the theorem tells us that exactly 760 Claude-like decompositions are valid for all odd $m > 1$.

Maybe the decomposition that Claude found wasn’t the “nicest” of those 760. Can we do better? I looked at several of them and found that they have somewhat different behaviors. Yet I didn’t encounter any that were actually nicer.

The dependence on i , j , and s means that these decompositions don’t have cyclic symmetry. I did notice, to my surprise, that 136 of the 760 generalizable $3 \times 3 \times 3$ cycles remain generalizable when we map ijk to jki . However, none are common to all three mappings $\{ijk, jki, kij\}$.

Filip told me that the explorations reported above, though ultimately successful, weren’t really smooth. He had to do some restarts when Claude stopped on random errors; then some of the previous search results were lost. After every two or three test programs were run, he had to remind Claude again and again that it was supposed to document its progress carefully.

Delicious success for odd m , at exploration number 31, came about one hour after the session began.

This decomposition problem remains open for even values of m . The case $m = 2$ was proved impossible long ago [1]. As part of exploration number 24, Claude said that it had found solutions for $m = 4$, $m = 6$, and $m = 8$; yet it saw no way to generalize those results.

Filip also told me that he asked Claude to continue on the even case after the odd case had been resolved. “But there after a while it seemed to get stuck. In the end, it was not even able to write and run explore programs correctly anymore, very weird. So I stopped the search.”

All in all, however, this was definitely an impressive success story. I think Claude Shannon’s spirit is probably proud to know that his name is now being associated with such advances. Hats off to Claude!

Appendix. Claude’s second cycle ($c = 1$) is governed by the following rules: “If $s = 0$, bump j . If $0 < s < m+1$, bump i . If $s = m-1$ and $i > 0$, bump k . If $s = m-1$ and $i = 0$, bump j .”

We can show that the vertices with $s = 0$ are seen in the following order, for $k = 0, 1, \dots, m-1$: $0k(-k)$, $(-2)(1+k)(1-k)$, $(-4)(2+k)(2-k)$, \dots , $2(-1+k)(-1-k)$. And that will establish the order in which vertices with $s = 1, 2, \dots, m-1$ are seen.

Claude’s third cycle ($c = 2$) is governed by somewhat more complex rules: “If $s = 0$ and $j < m-1$, bump i . If $s = 0$ and $j = m-1$, bump k . If $0 < s < m-1$ and $i < m-1$, bump k . If $0 < s < m-1$ and $i = m-1$, bump j . If $s = m-1$, bump i .”

The vertices with $s = 0$ and $j < m-1$ are seen in the following order: $0j(-j)$, $2j(-2-j)$, $4j(-4-j)$, \dots , $(-2)j(2-j)$. And the successors of the latter vertex are $(-1)j(2-j)$, $(-1)(j+1)(2-j)$, $(-1)(j+2)(2-j)$, \dots , $(-1)(j-2)(2-j)$, $0(j-2)(2-j)$.

But the vertices with $s = 0$ and $j = m-1$ are seen thus: $0(-1)1$, $1(-1)0$, $2(-1)(-1)$, \dots , $(-1)(-1)2$. And the successors of the latter vertex are $(-1)(-1)3$, $(-1)03$, $(-1)13$, \dots , $(-1)(-3)3$, $0(-3)3$.

Thus in every case the sequence of m vertices for $s = 0$ and a given j is followed by the sequence for $s = 0$ and $j-2$, modulo m .

References.

- [1] Jacques Aubert and Bernadette Schneider, “Graphes orientes indécomposables en circuits hamiltoniens.” *Journal of Combinatorial Theory* **B32** (1982), 347–349.
- [2] Donald E. Knuth, *The Art of Computer Programming*, Volume 4A: *Combinatorial Algorithms, Part 1* (Upper Saddle River, N. J.: Addison–Wesley, 2011), xvi+883 pp.
- [3] Donald E. Knuth, preliminary drafts entitled “Hamiltonian paths and cycles,” currently posted at <https://cs.stanford.edu/~knuth/fasc8a.ps.gz> and updated frequently as more material is being accumulated.